

MAO-A Polyclonal Antibody

Catalog # AP73888

Product Information

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | P21397 |
| Reactivity | Human, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 59682 |

Additional Information

| | |
|--------------------|---|
| Gene ID | 4128 |
| Other Names | MAOA; Amine oxidase [flavin-containing] A; Monoamine oxidase type A; MAO-A |
| Dilution | WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. |
| Format | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide. |
| Storage Conditions | -20°C |


Protein Information

| | |
|-------------------|--|
| Name | MAOA (HGNC:6833) |
| Function | Catalyzes the oxidative deamination of primary and some secondary amine such as neurotransmitters, with concomitant reduction of oxygen to hydrogen peroxide and has important functions in the metabolism of neuroactive and vasoactive amines in the central nervous system and peripheral tissues (PubMed: 18391214 , PubMed: 20493079 , PubMed: 24169519 , PubMed: 8316221). Preferentially oxidizes serotonin (PubMed: 20493079 , PubMed: 24169519). Also catalyzes the oxidative deamination of kynuramine to 3-(2-aminophenyl)-3-oxopropanal that can spontaneously condense to 4-hydroxyquinoline (By similarity). |
| Cellular Location | Mitochondrion outer membrane {ECO:0000250 UniProtKB:P21396}; Single-pass type IV membrane protein {ECO:0000250 UniProtKB:P21396}; Cytoplasmic side {ECO:0000250 UniProtKB:P21396} |
| Tissue Location | Heart, liver, duodenum, blood vessels and kidney. |

Background

Catalyzes the oxidative deamination of biogenic and xenobiotic amines and has important functions in the metabolism of neuroactive and vasoactive amines in the central nervous system and peripheral tissues. MAOA preferentially oxidizes biogenic amines such as 5-hydroxytryptamine (5-HT), norepinephrine and epinephrine.

Images



Western blot analysis of 293t lysis using MAOA antibody. Antibody was diluted at 1:500. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.