

# **IL-22 Polyclonal Antibody**

Catalog # AP73905

#### **Product Information**

Application WB, E
Primary Accession Q9GZX6

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW20011

### **Additional Information**

**Gene ID** 50616

Other Names IL22; ILTIF; ZCYTO18; Interleukin-22; IL-22; Cytokine Zcyto18; IL-10-related

T-cell-derived-inducible factor; IL-TIF

Dilution WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications. E~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

#### **Protein Information**

Name IL22

Synonyms ILTIF, ZCYTO18

**Function** Cytokine that plays a critical role in modulating tissue responses during inflammation (PubMed: 17204547). Plays an essential role in the regeneration

of epithelial cells to maintain barrier function after injury and for the prevention of further tissue damage (PubMed:17204547). Unlike most of the cytokines, has no effect on immune cells. Signals through a heterodimeric receptor composed of two subunits, the specific receptor IL22RA1 which is present on non-immune cells in many organs and the shared subunit IL10RB (PubMed:10875937, PubMed:18599299). Ligation of IL22RA1 with IL22 induces activation of the tyrosine kinases JAK1 and TYK2, which in turn

STAT3, ERK1/2 and PI3K/AKT pathways (PubMed: 25793261,

PubMed:31311100). Promotes phosphorylation of GSK3B at 'Ser-9' and CTTN

activates STAT3. In turn, promotes cell survival and proliferation through


(By similarity). Promotes epithelial cell spreading (By similarity).

**Cellular Location** Secreted.

# **Background**

Cytokine that contributes to the inflammatory response in vivo.

## **Images**



Western blot analysis of MOUSE-KIDNEY lysis using IL22 antibody. Antibody was diluted at 1:500. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.