

IL-22 Polyclonal Antibody

Catalog # AP73905

Product Information

Application WB, E
Primary Accession Q9GZX6

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW20011

Additional Information

Gene ID 50616

Other Names IL22; ILTIF; ZCYTO18; Interleukin-22; IL-22; Cytokine Zcyto18; IL-10-related

T-cell-derived-inducible factor; IL-TIF

Dilution WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications. E~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name IL22

Synonyms ILTIF, ZCYTO18

Function Cytokine that plays a critical role in modulating tissue responses during inflammation (PubMed: 17204547). Plays an essential role in the regeneration

of epithelial cells to maintain barrier function after injury and for the prevention of further tissue damage (PubMed:<u>17204547</u>). Unlike most of the cytokines, has no effect on immune cells. Signals through a heterodimeric receptor composed of two subunits, the specific receptor IL22RA1 which is present on non-immune cells in many organs and the shared subunit IL10RB (PubMed:<u>10875937</u>, PubMed:<u>18599299</u>). Ligation of IL22RA1 with IL22 induces activation of the tyrosine kinases JAK1 and TYK2, which in turn

STAT3, ERK1/2 and PI3K/AKT pathways (PubMed: 25793261,

PubMed:31311100). Promotes phosphorylation of GSK3B at 'Ser-9' and CTTN

activates STAT3. In turn, promotes cell survival and proliferation through


(By similarity). Promotes epithelial cell spreading (By similarity).

Cellular Location Secreted.

Background

Cytokine that contributes to the inflammatory response in vivo.

Images



Western blot analysis of MOUSE-KIDNEY lysis using IL22 antibody. Antibody was diluted at 1:500. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.