

CLECSF6 Polyclonal Antibody

Catalog # AP73990

Product Information

Application	WB
Primary Accession	Q9UMR7
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27512

Additional Information

Gene ID	50856
Other Names	C-type lectin domain family 4 member A (C-type lectin DDB27) (C-type lectin superfamily member 6) (Dendritic cell immunoreceptor) (Lectin-like immunoreceptor)
Dilution	WB~~WB 1:500-2000, ELISA 1:10000-20000
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	CLEC4A (HGNC:13257)
Function	C-type lectin receptor that binds carbohydrates mannose and fucose but also weakly interacts with N-acetylglucosamine (GlcNAc) in a Ca(2+)-dependent manner (PubMed: 27015765). Involved in regulating immune reactivity (PubMed: 10438934 , PubMed: 18258799). Once triggered by antigen, it is internalized by clathrin-dependent endocytosis and delivers its antigenic cargo into the antigen presentation pathway resulting in cross-priming of CD8(+) T cells. This cross-presentation and cross-priming are enhanced by TLR7 and TLR8 agonists with increased expansion of the CD8(+) T cells, high production of IFNG and TNF with reduced levels of IL4, IL5 and IL13 (PubMed: 18258799 , PubMed: 20530286). In plasmacytoid dendritic cells, inhibits TLR9-mediated IFNA and TNF production (PubMed: 18258799). May be involved via its ITIM motif (immunoreceptor tyrosine-based inhibitory motifs) in the inhibition of B-cell-receptor-mediated calcium mobilization and protein tyrosine phosphorylation (PubMed: 10438934).
Cellular Location	Cell membrane; Single-pass type II membrane protein; Extracellular side


Tissue Location

Expressed preferentially in hematopoietic tissues. Expressed in all circulating Ag-presenting cells such as dendritic cells, myeloid cells, monocytes, macrophages, B-cells and epidermal Langerhans cells (at protein level). Expressed in peripheral blood leukocytes, neutrophils, moderate quantities in spleen, lymph node, and bone marrow, and at very low levels in thymus


Background

C-type lectin receptor that binds carbohydrates mannose and fucose but also weakly interacts with N-acetylglucosamine (GlcNAc) in a Ca(2+)-dependent manner (PubMed:[27015765](#)). Involved in regulating immune reactivity (PubMed:[18258799](#), PubMed:[10438934](#)). Once triggered by antigen, it is internalized by clathrin-dependent endocytosis and delivers its antigenic cargo into the antigen presentation pathway resulting in cross-priming of CD8(+) T cells. This cross-presentation and cross-priming are enhanced by TLR7 and TLR8 agonists with increased expansion of the CD8(+) T cells, high production of IFNG and TNF with reduced levels of IL4, IL5 and IL13 (PubMed:[18258799](#), PubMed:[20530286](#)). In plasmacytoid dendritic cells, inhibits TLR9-mediated IFNA and TNF production (PubMed:[18258799](#)). May be involved via its ITIM motif (immunoreceptor tyrosine-based inhibitory motifs) in the inhibition of B-cell-receptor-mediated calcium mobilization and protein tyrosine phosphorylation (PubMed:[10438934](#)).


Images



Western Blot analysis of HEPG2 cells using CLECSF6 Polyclonal Antibody diluted at 1:500. Secondary antibody was diluted at 1:20000





Immunohistochemical analysis of paraffin-embedded human-colon-cancer, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-colon-cancer, antibody was diluted at 1:200

Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.