

# E2F-1 Polyclonal Antibody

Catalog # AP73994

## Product Information

Application	WB, IHC-P, IF
Primary Accession	<a href="#">Q01094</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46920

## Additional Information

Gene ID	1869
Other Names	E2F1 RBBP3
Dilution	WB~~IF: 1:50-200 WB 1:500-2000, ELISA 1:10000-20000 IHC 1:50-300 IHC-P~~IF: 1:50-200 WB 1:500-2000, ELISA 1:10000-20000 IHC 1:50-300 IF~~IF: 1:50-200 WB 1:500-2000, ELISA 1:10000-20000 IHC 1:50-300
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


## Protein Information

Name	E2F1 {ECO:0000303 PubMed:8964493, ECO:0000312 HGNC:HGNC:3113}
Function	<p>Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication (PubMed:<a href="#">10675335</a>, PubMed:<a href="#">12717439</a>, PubMed:<a href="#">17050006</a>, PubMed:<a href="#">17704056</a>, PubMed:<a href="#">18625225</a>, PubMed:<a href="#">28992046</a>). The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase (PubMed:<a href="#">10675335</a>, PubMed:<a href="#">12717439</a>, PubMed:<a href="#">17704056</a>). E2F1 binds preferentially RB1 in a cell-cycle dependent manner (PubMed:<a href="#">10675335</a>, PubMed:<a href="#">12717439</a>, PubMed:<a href="#">17704056</a>). It can mediate both cell proliferation and TP53/p53-dependent apoptosis (PubMed:<a href="#">8170954</a>). Blocks adipocyte differentiation by binding to specific promoters repressing CEBPA binding to its target gene promoters (PubMed:<a href="#">20176812</a>). Directly activates transcription of PEG10 (PubMed:<a href="#">17050006</a>, PubMed:<a href="#">18625225</a>, PubMed:<a href="#">28992046</a>). Positively regulates transcription of RRP1B (PubMed:<a href="#">20040599</a>).</p>
Cellular Location	Nucleus


## Background

Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC- 3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication. The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase. E2F1 binds preferentially RB1 in a cell-cycle dependent manner. It can mediate both cell proliferation and TP53/p53-dependent apoptosis. Blocks adipocyte differentiation by binding to specific promoters repressing CEBPA binding to its target gene promoters (PubMed:[20176812](#)). Positively regulates transcription of RRP1B (PubMed:[20040599](#)).


## Images




Immunofluorescence analysis of rat-heart tissue. 1,E2F-1 Polyclonal Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B



Immunofluorescence analysis of rat-heart tissue. 1,E2F-1 Polyclonal Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B



Immunohistochemical analysis of paraffin-embedded Mouse-lung tissue. 1,E2F-1 Polyclonal Antibody was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.



Western Blot analysis of MOUSE-BRAIN cells using Antibody diluted at 500. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.