

CACTIN Polyclonal Antibody

Catalog # AP74009

Product Information

Application	WB
Primary Accession	Q8WUQ7
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	88702

Additional Information

Gene ID	58509
Other Names	CACTIN C19orf29
Dilution	WB~~WB 1:500-2000, ELISA 1:10000-20000
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	CACTIN
Synonyms	C19orf29
Function	Plays a role in pre-mRNA splicing by facilitating excision of a subset of introns (PubMed: 28062851). Required for the splicing of CDCA5/Sororin, a regulator of sister chromatid cohesion (PubMed: 28062851). Involved in the regulation of innate immune response (PubMed: 20829348). Acts as a negative regulator of Toll-like receptor, interferon-regulatory factor (IRF) and canonical NF-kappa-B signaling pathways (PubMed: 20829348 , PubMed: 26363554). Contributes to the regulation of transcriptional activation of NF-kappa-B target genes in response to endogenous pro-inflammatory stimuli (PubMed: 20829348 , PubMed: 26363554).
Cellular Location	Nucleus. Cytoplasm, cytosol. Note=Nuclear localization with a speckled expression pattern in some cells. Colocalizes with NFKBIL1 in the nucleus.

Background

Involved in the regulation of innate immune response. Acts as negative regulator of Toll-like receptor and

interferon- regulatory factor (IRF) signaling pathways. Contributes to the regulation of transcriptional activation of NF-kappa-B target genes in response to endogenous proinflammatory stimuli. May play a role during early embryonic development. Probably involved in pre-mRNA splicing.

Images



Western Blot analysis of hepg2 cells using Antibody diluted at 1000. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.