

IL-31 Polyclonal Antibody

Catalog # AP74024

Product Information

Application	WB
Primary Accession	Q6EBC2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	18205

Additional Information


Gene ID	386653
Other Names	Interleukin-31 (IL-31)
Dilution	WB~~WB 1:500-2000, ELISA 1:10000-20000
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	IL31
Function	Activates STAT3 and possibly STAT1 and STAT5 through the IL31 heterodimeric receptor composed of IL31RA and OSMR (PubMed: 15184896). May function in skin immunity (PubMed: 15184896). Enhances myeloid progenitor cell survival in vitro (By similarity). Induces RETNLA and serum amyloid A protein expression in macrophages (By similarity).
Cellular Location	Secreted.
Tissue Location	Detected at low levels in testis, bone marrow, skeletal muscle, kidney, colon, thymus, small intestine and trachea

Background


Activates STAT3 and possibly STAT1 and STAT5 through the IL31 heterodimeric receptor composed of IL31RA and OSMR (PubMed:[15184896](#)). May function in skin immunity (PubMed:[15184896](#)). Enhances myeloid progenitor cell survival in vitro (By similarity). Induces RETNLA and serum amyloid A protein expression in macrophages (By similarity).




Western blot analysis of Hela Cell Lysate using antibody.
Secondary antibody was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded
human-testis, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded
human-testis, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded
human-spleen, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded
human-spleen, antibody was diluted at 1:200

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.