

Parathyroid hormone/parathyroid hormone-related peptide receptor Polyclonal Antibody

Catalog # AP74039

Product Information

Application	WB, IHC-P
Primary Accession	Q03431
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	66361

Additional Information

Gene ID	5745
Other Names	Parathyroid hormone/parathyroid hormone-related peptide receptor (PTH/PTHrP type I receptor) (PTH/PTHr receptor) (Parathyroid hormone 1 receptor) (PTH1 receptor)
Dilution	WB~~WB 1:500-2000,IHC-p 1:500-200, ELISA 1:10000-20000 IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	PTH1R {ECO:0000303 PubMed:10913300, ECO:0000312 HGNC:HGNC:9608}
Function	G-protein-coupled receptor for parathyroid hormone (PTH) and for parathyroid hormone-related peptide (PTHrP) (PubMed: 10913300 , PubMed: 18375760 , PubMed: 19674967 , PubMed: 27160269 , PubMed: 30975883 , PubMed: 35932760 , PubMed: 8397094). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors, such as adenylate cyclase (cAMP) (PubMed: 30975883 , PubMed: 35932760). PTH1R is coupled to G(s) G alpha proteins and mediates activation of adenylate cyclase activity (PubMed: 20172855 , PubMed: 30975883 , PubMed: 35932760). PTHrP dissociates from PTH1R more rapidly than PTH; as consequence, the cAMP response induced by PTHrP decays faster than the response induced by PTH (PubMed: 35932760).
Cellular Location	Cell membrane; Multi-pass membrane protein


Tissue Location

Expressed in most tissues. Most abundant in kidney, bone and liver.


Background

Receptor for parathyroid hormone and for parathyroid hormone-related peptide. The activity of this receptor is mediated by G proteins which activate adenylyl cyclase and also a phosphatidylinositol-calcium second messenger system.


Images



Western blot analysis of MCF7 Cell Lysate, antibody was diluted at 1:1000. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.