10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Keratin-pan Polyclonal Antibody

Catalog # AP74053

Product Information

Application	WB, E, IHC-P
Primary Accession	P35908, Q01546, P12035, P13647, P02538, P04259, P48668, Q3SY84, Q14CN4, Q86Y46, Q7RTS7, O95678, Q5XKE5, P08729, P0
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	65433


Additional Information

Gene ID	3849
Dilution	WB~~WB 1:500-2000, ELISA 1:10000-20000 E~~N/A IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information

Name	KRT2 (HGNC:6439)
Synonyms	KRT2A, KRT2E
Function	Structural component of intermediate filaments in suprabasal keratinocytes of stratified epidermis. Forms heteropolymers with type I keratins (e.g., KRT10), assembling into keratin intermediate filament networks that support mechanical integrity and resilience of the differentiating epidermis (PubMed:12598329, PubMed:1380918). Contributes to terminal differentiation and cornification of keratinocytes and plays a role in establishing the epidermal barrier (PubMed:9804344).
Cellular Location	Cytoplasm.
Tissue Location	Expressed in the upper spinous and granular suprabasal layers of normal adult epidermal tissues from most body sites including thigh, breast nipple, foot sole, penile shaft and axilla. Not present in foreskin, squamous metaplasias and carcinomas Expression in hypertrophic and keloid scars begins in the deepest suprabasal layer. Weakly expressed in normal gingiva and tongue, however expression is induced in benign keratoses of lingual mucosa and in mild-to-moderate oral dysplasia with orthokeratinization

Images



Western blot analysis of mouse-lung mouse-brain mouse-kidney mouse-heart lysate, antibody was diluted at 500. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.