

# Keratin-pan Polyclonal Antibody

Catalog # AP74053

## Product Information

---

Application	WB
Primary Accession	<a href="#">P35908</a> , <a href="#">Q01546</a> , <a href="#">P12035</a> , <a href="#">P13647</a> , <a href="#">P02538</a> , <a href="#">P04259</a> , <a href="#">P48668</a> , <a href="#">Q3SY84</a> , <a href="#">Q14CN4</a> , <a href="#">Q86Y46</a> , <a href="#">Q7RTS7</a> , <a href="#">Q95678</a> , <a href="#">Q5XKE5</a> , <a href="#">P08729</a> , <a href="#">P0</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	65433

## Additional Information

---

Gene ID	3849
Dilution	WB~~WB 1:500-2000, ELISA 1:10000-20000
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


## Protein Information

---

Name	KRT2
Synonyms	KRT2A, KRT2E
Function	Probably contributes to terminal cornification (PubMed: <a href="#">1380918</a> ). Associated with keratinocyte activation, proliferation and keratinization (PubMed: <a href="#">12598329</a> ). Required for maintenance of corneocytes and keratin filaments in suprabasal keratinocytes in the epidermis of the ear, potentially via moderation of expression and localization of keratins and their partner proteins (By similarity). Plays a role in the establishment of the epidermal barrier on plantar skin (By similarity).
Cellular Location	Cytoplasm.
Tissue Location	Expressed in the upper spinous and granular suprabasal layers of normal adult epidermal tissues from most body sites including thigh, breast nipple, foot sole, penile shaft and axilla. Not present in foreskin, squamous metaplasias and carcinomas Expression in hypertrophic and keloid scars begins in the deepest suprabasal layer. Weakly expressed in normal gingiva and tongue, however expression is induced in benign keratoses of lingual mucosa and in mild-to-moderate oral dysplasia with orthokeratinization

## Images

---



Western blot analysis of mouse-lung mouse-brain mouse-kidney mouse-heart lysate, antibody was diluted at 500. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.