

Endoglin Polyclonal Antibody

Catalog # AP74062

Product Information

| | |
|--------------------------|------------------------|
| Application | WB, IHC-P, IF, ICC, E |
| Primary Accession | P17813 |
| Reactivity | Human, Rat, Mouse |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 70578 |

Additional Information

| | |
|---------------------------|---|
| Gene ID | 2022 |
| Other Names | Endoglin (CD antigen CD105) |
| Dilution | WB~~WB 1:500-2000,IHC-p 1:500-200, ELISA 1:10000-20000 IHC-P~~WB 1:500-2000,IHC-p 1:500-200, ELISA 1:10000-20000 IF~~1:50~200 ICC~~N/A E~~N/A |
| Format | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide. |
| Storage Conditions | -20°C |

Protein Information

| | |
|-----------------|--|
| Name | ENG |
| Synonyms | END |
| Function | Vascular endothelium glycoprotein that plays an important role in the regulation of angiogenesis (PubMed: 21737454 , PubMed: 23300529). Required for normal structure and integrity of adult vasculature (PubMed: 7894484). Regulates the migration of vascular endothelial cells (PubMed: 17540773). Required for normal extraembryonic angiogenesis and for embryonic heart development (By similarity). May regulate endothelial cell shape changes in response to blood flow, which drive vascular remodeling and establishment of normal vascular morphology during angiogenesis (By similarity). May play a critical role in the binding of endothelial cells to integrins and/or other RGD receptors (PubMed: 1692830). Acts as a TGF-beta coreceptor and is involved in the TGF-beta/BMP signaling cascade that ultimately leads to the activation of SMAD transcription factors (PubMed: 21737454 , PubMed: 22347366 , PubMed: 23300529 , PubMed: 8370410). Required for GDF2/BMP9 signaling through SMAD1 in endothelial cells and modulates TGFB1 signaling through SMAD3 (PubMed: 21737454 , PubMed: 22347366 , PubMed: 23300529). |

Cellular Location

Cell membrane; Single-pass type I membrane protein


Tissue Location

Detected on umbilical vein endothelial cells (PubMed:10625079). Detected in placenta (at protein level) (PubMed:1692830). Detected on endothelial cells (PubMed:1692830)

Background


Vascular endothelium glycoprotein that plays an important role in the regulation of angiogenesis (PubMed:[21737454](#), PubMed:[23300529](#)). Required for normal structure and integrity of adult vasculature (PubMed:[7894484](#)). Regulates the migration of vascular endothelial cells (PubMed:[17540773](#)). Required for normal extraembryonic angiogenesis and for embryonic heart development (By similarity). May regulate endothelial cell shape changes in response to blood flow, which drive vascular remodeling and establishment of normal vascular morphology during angiogenesis (By similarity). May play a critical role in the binding of endothelial cells to integrins and/or other RGD receptors (PubMed:[1692830](#)). Acts as TGF-beta coreceptor and is involved in the TGF-beta/BMP signaling cascade that ultimately leads to the activation of SMAD transcription factors (PubMed:[8370410](#), PubMed:[21737454](#), PubMed:[22347366](#), PubMed:[23300529](#)). Required for GDF2/BMP9 signaling through SMAD1 in endothelial cells and modulates TGF β 1 signaling through SMAD3 (PubMed:[21737454](#), PubMed:[22347366](#), PubMed:[23300529](#)).

Images




Western blot analysis of 3T3 KB K562 HeLa 293T lysate, antibody was diluted at 500. Secondary antibody was diluted at 1:20000


15--




Immunohistochemical analysis of paraffin-embedded human-tonsil, antibody was diluted at 1:200




Immunohistochemical analysis of paraffin-embedded human-tonsil, antibody was diluted at 1:200




Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200




Immunohistochemical analysis of paraffin-embedded Human Amygdala. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human Amygdala. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).





Immunohistochemical analysis of paraffin-embedded Human Amygdala. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.