

# IFN- $\alpha$ / $\beta$ R $\alpha$ Polyclonal Antibody

Catalog # AP74089

## Product Information

---

Application	IHC-P
Primary Accession	<a href="#">P17181</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	63525

## Additional Information

---

Gene ID	3454
Other Names	Interferon alpha/beta receptor 1 (IFN-R-1) (IFN-alpha/beta receptor 1) (Cytokine receptor class-II member 1) (Cytokine receptor family 2 member 1) (CRF2-1) (Type I interferon receptor 1)
Dilution	IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

---

Name	IFNAR1
Synonyms	IFNAR
Function	<p>Together with IFNAR2, forms the heterodimeric receptor for type I interferons (including interferons alpha, beta, epsilon, omega and kappa) (PubMed:<a href="#">10049744</a>, PubMed:<a href="#">14532120</a>, PubMed:<a href="#">15337770</a>, PubMed:<a href="#">2153461</a>, PubMed:<a href="#">21854986</a>, PubMed:<a href="#">24075985</a>, PubMed:<a href="#">31270247</a>, PubMed:<a href="#">33252644</a>, PubMed:<a href="#">35442418</a>, PubMed:<a href="#">7813427</a>). Type I interferon binding activates the JAK-STAT signaling cascade, resulting in transcriptional activation or repression of interferon-regulated genes that encode the effectors of the interferon response (PubMed:<a href="#">10049744</a>, PubMed:<a href="#">21854986</a>, PubMed:<a href="#">7665574</a>). Mechanistically, type I interferon- binding brings the IFNAR1 and IFNAR2 subunits into close proximity with one another, driving their associated Janus kinases (JAKs) (TYK2 bound to IFNAR1 and JAK1 bound to IFNAR2) to cross-phosphorylate one another (PubMed:<a href="#">21854986</a>, PubMed:<a href="#">32972995</a>, PubMed:<a href="#">7665574</a>, PubMed:<a href="#">7813427</a>). The activated kinases phosphorylate specific tyrosine residues on the intracellular domains of IFNAR1 and IFNAR2, forming docking sites for the STAT transcription</p>

factors (PubMed:[21854986](#), PubMed:[32972995](#), PubMed:[7526154](#), PubMed:[7665574](#), PubMed:[7813427](#)). STAT proteins are then phosphorylated by the JAKs, promoting their translocation into the nucleus to regulate expression of interferon-regulated genes (PubMed:[19561067](#), PubMed:[21854986](#), PubMed:[32972995](#), PubMed:[7665574](#), PubMed:[7813427](#), PubMed:[9121453](#)). Can also act independently of IFNAR2: form an active IFNB1 receptor by itself and activate a signaling cascade that does not involve activation of the JAK-STAT pathway (By similarity).

#### Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein. Late endosome. Lysosome. Note=Interferon binding triggers internalization of the receptor from the cell membrane into endosomes and then into lysosomes.

#### Tissue Location

IFN receptors are present in all tissues and even on the surface of most IFN-resistant cells. Isoform 1, isoform 2 and isoform 3 are expressed in the IFN-alpha sensitive myeloma cell line U266B1. Isoform 2 and isoform 3 are expressed in the IFN-alpha resistant myeloma cell line U266R. Isoform 1 is not expressed in IFN- alpha resistant myeloma cell line U266R.


## Background

---


Component of the receptor for type I interferons, including interferons alpha, IFNB1 and IFNW1 (PubMed:[2153461](#), PubMed:[7665574](#), PubMed:[10049744](#), PubMed:[14532120](#), PubMed:[15337770](#), PubMed:[21854986](#)). Functions in general as heterodimer with IFNAR2 (PubMed:[7665574](#), PubMed:[10049744](#), PubMed:[21854986](#)). Type I interferon binding activates the JAK-STAT signaling cascade, and triggers tyrosine phosphorylation of a number of proteins including JAKs, TYK2, STAT proteins and the IFNR alpha- and beta- subunits themselves (PubMed:[7665574](#), PubMed:[21854986](#)). Can form an active IFNB1 receptor by itself and activate a signaling cascade that does not involve activation of the JAK-STAT pathway (By similarity).

## Images

---



Immunohistochemical analysis of paraffin-embedded human-lung-cancer, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-lung-cancer, antibody was diluted at 1:200