

CD204 Polyclonal Antibody

Catalog # AP74130

Product Information

Application	IHC-P, IF, ICC, E
Primary Accession	P21757
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	49762

Additional Information

Gene ID	4481
Other Names	Macrophage scavenger receptor types I and II (Macrophage acetylated LDL receptor I and II) (Scavenger receptor class A member 1) (CD antigen CD204)
Dilution	IHC-P~~IHC-p 1:50-200, ELISA 1:10000-20000 IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	MSR1
Synonyms	SCARA1
Function	Membrane glycoproteins implicated in the pathologic deposition of cholesterol in arterial walls during atherogenesis. Two types of receptor subunits exist. These receptors mediate the endocytosis of a diverse group of macromolecules, including modified low density lipoproteins (LDL) (PubMed: 2251254). Isoform III does not internalize acetylated LDL (PubMed: 9548586).
Cellular Location	Membrane; Single-pass type II membrane protein.
Tissue Location	Isoform I, isoform II and isoform III are expressed in monocyte-derived macrophages. Isoform I and isoform II are expressed in the liver, placenta and brain.


Background

Membrane glycoproteins implicated in the pathologic deposition of cholesterol in arterial walls during atherogenesis. Two types of receptor subunits exist. These receptors mediate the endocytosis of a diverse group of macromolecules, including modified low density lipoproteins (LDL) (PubMed:[2251254](#)). Isoform III does not internalize acetylated LDL (PubMed:[9548586](#)).

Images



Immunohistochemical analysis of paraffin-embedded human liver, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human liver, antibody was diluted at 1:200

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.