

VHL Polyclonal Antibody

Catalog # AP74171

Product Information

Application	IHC-P
Primary Accession	P40337
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24153

Additional Information

Gene ID	7428
Other Names	Von Hippel-Lindau disease tumor suppressor (Protein G7) (pVHL)
Dilution	IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	VHL
Function	Involved in the ubiquitination and subsequent proteasomal degradation via the von Hippel-Lindau ubiquitination complex (PubMed: 10944113 , PubMed: 17981124 , PubMed: 19584355). Seems to act as a target recruitment subunit in the E3 ubiquitin ligase complex and recruits hydroxylated hypoxia-inducible factor (HIF) under normoxic conditions (PubMed: 10944113 , PubMed: 17981124). Involved in transcriptional repression through interaction with HIF1A, HIF1AN and histone deacetylases (PubMed: 10944113 , PubMed: 17981124). Ubiquitinates, in an oxygen-responsive manner, ADRB2 (PubMed: 19584355). Acts as a negative regulator of mTORC1 by promoting ubiquitination and degradation of RPTOR (PubMed: 34290272).
Cellular Location	[Isoform 1]: Cytoplasm. Cell membrane; Peripheral membrane protein. Endoplasmic reticulum. Nucleus. Note=Found predominantly in the cytoplasm and with less amounts nuclear or membrane-associated (PubMed:9751722) Colocalizes with ADRB2 at the cell membrane (PubMed:19584355)
Tissue Location	Expressed in the adult and fetal brain and kidney.


Background

Involved in the ubiquitination and subsequent proteasomal degradation via the von Hippel-Lindau ubiquitination complex. Seems to act as a target recruitment subunit in the E3 ubiquitin ligase complex and recruits hydroxylated hypoxia-inducible factor (HIF) under normoxic conditions. Involved in transcriptional repression through interaction with HIF1A, HIF1AN and histone deacetylases. Ubiquitinates, in an oxygen-responsive manner, ADRB2.

Images



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:200

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.