

ST2 Polyclonal Antibody

Catalog # AP74175

Product Information

Application IHC-P
Primary Accession Q01638
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 63358

Additional Information

Gene ID 9173

Other Names Interleukin-1 receptor-like 1 (Protein ST2)

Dilution IHC-P~~IHC-p 1:50-200, ELISA 1:10000-20000

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name IL1RL1

Synonyms DER4, ST2, T1

Function Receptor for interleukin-33 (IL-33) which plays crucial roles in innate and

adaptive immunity, contributing to tissue homeostasis and responses to environmental stresses together with coreceptor IL1RAP (PubMed:35238669). Its stimulation recruits MYD88, IRAK1, IRAK4, and TRAF6, followed by

phosphorylation of MAPK3/ERK1 and/or MAPK1/ERK2, MAPK14, and MAPK8. Possibly involved in helper T-cell function (Probable) (PubMed:16286016). Upon tissue injury, induces UCP2- dependent mitochondrial rewiring that attenuates the generation of reactive oxygen species and preserves the integrity of Krebs cycle required for persistent production of itaconate and subsequent GATA3- dependent differentiation of inflammation-resolving

alternatively activated macrophages (By similarity).

Cellular Location [Isoform C]: Cell membrane. Cell membrane; Single-pass type I membrane

protein

Tissue Location Highly expressed in kidney, lung, placenta, stomach, skeletal muscle, colon


and small intestine. Isoform A is prevalently expressed in the lung, testis,

placenta, stomach and colon Isoform B is more abundant in the brain, kidney and the liver. Isoform C is not detected in brain, heart, liver, kidney and skeletal muscle Expressed on T-cells in fibrotic liver; at protein level. Overexpressed in fibrotic and cirrhotic liver.

Background

Receptor for interleukin-33 (IL-33); signaling requires association of the coreceptor IL1RAP. Its stimulation recruits MYD88, IRAK1, IRAK4, and TRAF6, followed by phosphorylation of MAPK3/ERK1 and/or MAPK1/ERK2, MAPK14, and MAPK8. Possibly involved in helper T-cell function.

Images



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.