

Noggin Polyclonal Antibody

Catalog # AP74176

Product Information

Application IHC-P, IF, ICC, E

Primary Accession Q13253

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW25774

Additional Information

Gene ID 9241

Other Names Noggin

Dilution IHC-P~~IHC-p 1:50-200, ELISA 1:10000-20000 IF~~1:50~200 ICC~~N/A E~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name NOG

Function Inhibitor of bone morphogenetic proteins (BMP) signaling which is required

for growth and patterning of the neural tube and somite. Essential for cartilage morphogenesis and joint formation. Inhibits chondrocyte differentiation through its interaction with GDF5 and, probably, GDF6


(PubMed: 21976273, PubMed: 26643732).

Cellular Location Secreted.

Background

Inhibitor of bone morphogenetic proteins (BMP) signaling which is required for growth and patterning of the neural tube and somite. Essential for cartilage morphogenesis and joint formation. Inhibits chondrocyte differentiation through its interaction with GDF5 and, probably, GDF6 (PubMed:21976273, PubMed:26643732).

Images



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.