

Cytoglobin Polyclonal Antibody

Catalog # AP74205

Product Information

Application	IHC-P, IF, ICC, E
Primary Accession	Q8WWM9
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21405

Additional Information

Gene ID	114757
Other Names	Cytoglobin (Histoglobin) (HGb) (Stellate cell activation-associated protein)
Dilution	IHC-P~~IHC-p 1:50-200, ELISA 1:10000-20000 IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	CYGB (HGNC:16505)
Function	Probable multifunctional globin with a hexacoordinated heme iron required for the catalysis of various reactions depending on redox condition of the cell as well as oxygen availability (PubMed: 11893755 , PubMed: 12359339 , PubMed: 15165856 , PubMed: 19147491 , PubMed: 20511233 , PubMed: 28393874 , PubMed: 28671819 , PubMed: 29128400 , PubMed: 33576020 , PubMed: 34930834). Has a nitric oxide dioxygenase (NOD) activity and is most probably involved in cell-mediated and oxygen-dependent nitric oxide consumption (PubMed: 19147491 , PubMed: 20511233 , PubMed: 28393874 , PubMed: 28671819). By scavenging this second messenger may regulate several biological processes including endothelium-mediated vasodilation and vascular tone (PubMed: 19147491 , PubMed: 28393874). Under normoxic conditions functions as a nitric oxide dioxygenase (NOD) but under hypoxic conditions the globin may switch its function to that of a nitrite (NO ₂) reductase (NiR), generating nitric oxide (PubMed: 29128400). Could also have peroxidase and superoxide dismutase activities, detoxifying reactive oxygen species and protecting cells against oxidative stress (PubMed: 12359339 , PubMed: 33576020 , PubMed: 34930834). Also binds dioxygen with low affinity and could function as an oxygen sensor but has probably no function as a respiratory oxygen carrier (PubMed: 11893755 ,

PubMed:[15299006](#), PubMed:[20553503](#)).

Cellular Location

Cytoplasm. Nucleus


Tissue Location

Widely expressed. Highest expression in heart, stomach, bladder and small intestine.


Background

May have a protective function during conditions of oxidative stress. May be involved in intracellular oxygen storage or transfer.


Images




Immunohistochemical analysis of paraffin-embedded Human-heart, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-heart, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-skeletal-muscle, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-skeletal-muscle, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.