

CD70 Polyclonal Antibody

Catalog # AP74214

Product Information

Application	IHC-P
Primary Accession	P32970
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21118

Additional Information

Gene ID	970
Other Names	CD70 antigen (CD27 ligand) (CD27-L) (Tumor necrosis factor ligand superfamily member 7) (CD antigen CD70)
Dilution	IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	CD70 (HGNC:11937)
Function	Expressed at the plasma membrane of B cells, it is the ligand of the CD27 receptor which is specifically expressed at the surface of T cells (PubMed: 28011863 , PubMed: 28011864 , PubMed: 8387892). The CD70- CD27 signaling pathway mediates antigen-specific T cell activation and expansion which in turn provides immune surveillance of B cells (PubMed: 28011863 , PubMed: 28011864).
Cellular Location	Cell membrane; Single-pass type II membrane protein


Background

Cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells.

Images



Immunohistochemical analysis of paraffin-embedded Human-tonsil, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-tonsil, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.