

# CD275 Polyclonal Antibody

Catalog # AP74221

## Product Information

---

Application	IHC-P
Primary Accession	<a href="#">O75144</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	33349

## Additional Information

---

Gene ID	23308
Other Names	ICOS ligand (B7 homolog 2) (B7-H2) (B7-like protein GI50) (B7-related protein 1) (B7RP-1) (CD antigen CD275)
Dilution	IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

---

Name	ICOSLG
Function	Ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell proliferation and cytokine secretion (PubMed: <a href="#">11007762</a> , PubMed: <a href="#">11023515</a> , PubMed: <a href="#">30498080</a> ). Also induces B-cell proliferation and differentiation into plasma cells. Could play an important role in mediating local tissue responses to inflammatory conditions, as well as in modulating the secondary immune response by co-stimulating memory T-cell function (By similarity). In endothelial cells, required for proper neutrophil transmigration in response to chemoattractants, such as CXCL8/IL8 or N-formyl-methionyl peptides (fMLP) (PubMed: <a href="#">30498080</a> ).
Cellular Location	Cell membrane; Single-pass type I membrane protein
Tissue Location	Expressed on peripheral blood B-cells and monocytes, as well as on monocyte-derived dendritic cells (at protein level). [Isoform 2]: Detected only in lymph nodes, leukocytes and spleen. Expressed on activated monocytes and dendritic cells.


## Background

---

Ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell proliferation and cytokine secretion; induces also B-cell proliferation and differentiation into plasma cells. Could play an important role in mediating local tissue responses to inflammatory conditions, as well as in modulating the secondary immune response by co- stimulating memory T-cell function (By similarity).

## Images

---



Immunohistochemical analysis of paraffin-embedded Human-tonsil, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.