10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Vinculin Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7426b

Product Information

Application	WB, E
Primary Accession	P18206
Other Accession	P85972, P26234, Q64727, P12003
Reactivity	Human, Mouse
Predicted	Rat, Pig, Chicken, Bovine, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB16690
Calculated MW	123799
Antigen Region	1012-1039

Additional Information

Gene ID	7414
Other Names	Vinculin, Metavinculin, MV, VCL
Target/Specificity	This VINC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1012-1039 amino acids from the C-terminal region of human VINC.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Vinculin Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	VCL
Function	Actin filament (F-actin)-binding protein involved in cell- matrix adhesion and cell-cell adhesion. Regulates cell-surface E- cadherin expression and potentiates mechanosensing by the E-cadherin complex. May also play

important roles in cell morphology and locomotion.

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:P12003}; Peripheral membrane protein {ECO:0000250|UniProtKB:P12003}; Cytoplasmic side {ECO:0000250|UniProtKB:P12003}. Cell junction, adherens junction {ECO:0000250|UniProtKB:P12003}. Cell junction, focal adhesion {ECO:0000250|UniProtKB:P12003}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P12003}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P85972}. Cell membrane, sarcolemma {ECO:0000250|UniProtKB:Q64727}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q64727}; Cytoplasmic side {ECO:0000250|UniProtKB:Q64727}. Cell projection, podosome {ECO:0000250|UniProtKB:Q64727}. Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:Q64727}. Note=Recruitment to cell-cell junctions occurs in a myosin II-dependent manner. Interaction with CTNNB1 is necessary for its localization to the cell-cell junctions {ECO:0000250|UniProtKB:P12003}

Tissue Location

Metavinculin is muscle-specific.


Background

VINC is a cytoskeletal protein associated with cell-cell and cell-matrix junctions, where it is thought to function as one of several interacting proteins involved in anchoring F-actin to the membrane. Defects in VCL are the cause of cardiomyopathy dilated type 1W. Dilated cardiomyopathy is a disorder characterized by ventricular dilation and impaired systolic function, resulting in congestive heart failure and arrhythmia.

References


Moiseyeva E.P., Weller P.A.J. Biol. Chem. 268:4318-4325(1993)
Sun N., Critchley D.R., Paulin D.Biochem. J. 409:657-667(2008)
Izard T., Evans G., Borgon R.A.Nature 427:171-175(2004)

Images



Anti-VINC Antibody (C-term) at 1:1000 dilution + Hela whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 124 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Western blot analysis of anti-Vinculin Antibody (C-term)(Cat.#AP7426b) in mouse lung tissue lysates (35ug/lane). VINC (arrow) was detected using the purified Pab.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.