

Tropomyosin α Polyclonal Antibody

Catalog # AP74272

Product Information

Application	WB
Primary Accession	P09493
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32709

Additional Information

Gene ID	7168
Other Names	Tropomyosin alpha-1 chain (Alpha-tropomyosin) (Tropomyosin-1)
Dilution	WB~~WB 1:500-2000, ELISA 1:10000-20000
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	TPM1
Synonyms	C15orf13, TMSA
Function	Binds to actin filaments in muscle and non-muscle cells (PubMed: 23170982). Plays a central role, in association with the troponin complex, in the calcium dependent regulation of vertebrate striated muscle contraction (PubMed: 23170982). Smooth muscle contraction is regulated by interaction with caldesmon. In non-muscle cells is implicated in stabilizing cytoskeleton actin filaments.
Cellular Location	Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:P04692}. Note=Associates with F-actin stress fibers. {ECO:0000250 UniProtKB:P04692}
Tissue Location	Detected in primary breast cancer tissues but undetectable in normal breast tissues in Sudanese patients. Isoform 1 is expressed in adult and fetal skeletal muscle and cardiac tissues, with higher expression levels in the cardiac tissues. Isoform 10 is expressed in adult and fetal cardiac tissues, but not in skeletal muscle. {ECO:0000269 PubMed:15249230, ECO:0000269 Ref.15}

Background

Binds to actin filaments in muscle and non-muscle cells (PubMed: [23170982](#)). Plays a central role, in association with the troponin complex, in the calcium dependent regulation of vertebrate striated muscle contraction (PubMed:[23170982](#)). Smooth muscle contraction is regulated by interaction with caldesmon. In non-muscle cells is implicated in stabilizing cytoskeleton actin filaments.

Images



Western Blot analysis of 1,mouse-lung 2,mouse-brain
3,mouse-spleen cells using primary antibody diluted at
1:500(4°C overnight). Secondary antibody : Goat
Anti-rabbit IgG IRDye 800(diluted at 1:5000, 25°C, 1 hour)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.