

TGF β1 Polyclonal Antibody

Catalog # AP74322

Product Information

Application WB Primary Accession P01137

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW44325

Additional Information

Gene ID 7040

Other Names Transforming growth factor beta-1 (TGF-beta-1) [Cleaved into:

Latency-associated peptide (LAP)]

Dilution WB~~WB 1:500-2000, ELISA 1:10000-20000

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name TGFB1 (HGNC:11766)

Synonyms TGFB

Function Transforming growth factor beta-1 proprotein: Precursor of the

Latency-associated peptide (LAP) and Transforming growth factor beta-1 (TGF-beta-1) chains, which constitute the regulatory and active subunit of

TGF-beta-1, respectively.

Cellular Location [Latency-associated peptide]: Secreted, extracellular space, extracellular

matrix

Tissue Location Highly expressed in bone (PubMed:11746498, PubMed:17827158).

Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes


with ASPN in chondrocytes within OA lesions of articular cartilage

(PubMed:17827158)

Background

Transforming growth factor beta-1 proprotein: Precursor of the Latency-associated peptide (LAP) and Transforming growth factor beta-1 (TGF-beta-1) chains, which constitute the regulatory and active subunit of TGF-beta-1, respectively.

Images



Western blot analysis of various lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.