

PAR1 (Cleaved-Ser42) Polyclonal Antibody

Catalog # AP74344

Product Information

Application	WB, E
Primary Accession	P25116
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47441

Additional Information

Gene ID	2149
Other Names	Proteinase-activated receptor 1 (PAR-1) (Coagulation factor II receptor) (Thrombin receptor)
Dilution	WB~~WB 1:500-2000, ELISA 1:10000-20000 E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	F2R (HGNC:3537)
Synonyms	CF2R, PAR1, TR
Function	High affinity receptor that binds the activated thrombin, leading to calcium release from intracellular stores (PubMed: 1672265 , PubMed: 8136362). The thrombin-activated receptor signaling pathway is mediated through PTX-insensitive G proteins, activation of phospholipase C resulting in the production of 1D-myo-inositol 1,4,5- trisphosphate (InsP3) which binds to InsP3 receptors causing calcium release from the stores (By similarity). In astrocytes, the calcium released into the cytosol allows the Ca(2+)-dependent release of L- glutamate into the synaptic cleft through BEST1, that targets the neuronal postsynaptic GRIN2A/NMDAR receptor resulting in the synaptic plasticity regulation (By similarity). May play a role in platelets activation and in vascular development (PubMed: 10079109). Mediates up- regulation of pro-inflammatory cytokines, such as MCP-1/CCL2 and IL6, triggered by coagulation factor Xa (F10) in cardiac fibroblasts and umbilical vein endothelial cells (PubMed: 30568593 , PubMed: 34831181).
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P26824}; Multi-pass membrane

protein {ECO:0000250|UniProtKB:P26824}


Tissue Location

Platelets and vascular endothelial cells.

Background

High affinity receptor for activated thrombin coupled to G proteins that stimulate phosphoinositide hydrolysis. May play a role in platelets activation and in vascular development.

Images



Western blot analysis of MCF-7 lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000

Citations

- [PAR1 regulation of CXCL1 expression and neutrophil recruitment to the lung in mice infected with influenza A virus](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.