

# ADAM9 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7437b

## Product Information

---

<b>Application</b>	WB, IHC-P, FC, E
<b>Primary Accession</b>	<a href="#">Q13443</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB17777
<b>Calculated MW</b>	90556
<b>Antigen Region</b>	704-733

## Additional Information

---

<b>Gene ID</b>	8754
<b>Other Names</b>	Disintegrin and metalloproteinase domain-containing protein 9, ADAM 9, 3424-, Cellular disintegrin-related protein, Meltrin-gamma, Metalloprotease/disintegrin/cysteine-rich protein 9, Myeloma cell metalloproteinase, ADAM9, KIAA0021, MCMP, MDC9, MLTNG
<b>Target/Specificity</b>	This ADAM9 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 704-733 amino acids from the C-terminal region of human ADAM9.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	ADAM9 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	ADAM9
<b>Synonyms</b>	KIAA0021, MCMP, MDC9, MLTNG

<b>Function</b>	Metalloprotease that cleaves and releases a number of molecules with important roles in tumorigenesis and angiogenesis, such as TEK, KDR, EPHB4, CD40, VCAM1 and CDH5. May mediate cell-cell, cell- matrix interactions and regulate the motility of cells via interactions with integrins.
<b>Cellular Location</b>	[Isoform 1]: Cell membrane; Single-pass type I membrane protein
<b>Tissue Location</b>	Widely expressed. Expressed in chondrocytes. Isoform 2 is highly expressed in liver and heart


## Background

ADAM9 is a member of the ADAM (a disintegrin and metalloprotease domain) family. Members of this family are membrane-anchored proteins structurally related to snake venom disintegrins, and have been implicated in a variety of biological processes involving cell-cell and cell-matrix interactions, including fertilization, muscle development, and neurogenesis. This protein interacts with SH3 domain-containing proteins, binds mitotic arrest deficient 2 beta protein, and is also involved in TPA-induced ectodomain shedding of membrane-anchored heparin-binding EGF-like growth factor.


## References

Weskamp G., Kraetzschmar J., Reid M.S.J. Cell Biol. 132:717-726(1996)  
Hotoda N., Koike H.Biochem. Biophys. Res. Commun. 293:800-805(2002)  
McKie N., Edwards T., Dallas D.J.Biochem. Biophys. Res. Commun. 230:335-339(1997)

## Images




Western blot analysis of ADAM9 Antibody (C-term) (Cat. #AP7437b) in MDA-MB231 cell line lysates (35ug/lane). ADAM9 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human lung carcinoma reacted with ADAM9 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

ADAM9 Antibody (C-term) (Cat. #AP7437b) flow cytometric analysis of MCF-7 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary



antibodies were used for the analysis.

## Citations

---

- [Loss of tumor suppressor miR-126 contributes to the development of hepatitis B virus-related hepatocellular carcinoma metastasis through the upregulation of ADAM9.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.