

CCL17 Polyclonal Antibody

Catalog # AP74470

Product Information

Application WB
Primary Accession Q92583
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 10507

Additional Information

Gene ID 6361

Other Names C-C motif chemokine 17 (CC chemokine TARC) (Small-inducible cytokine A17)

(Thymus and activation-regulated chemokine)

Dilution WB~~WB 1:500-2000, ELISA(peptide)1:5000-20000

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name CCL17

Synonyms SCYA17, TARC

Function Chemokine, which displays chemotactic activity for T lymphocytes,

preferentially Th2 cells, but not monocytes or granulocytes. Therefore plays an important role in a wide range of inflammatory and immunological processes (PubMed:8702936, PubMed:9169480). Acts by binding to CCR4 at

T-cell surface (PubMed: 10540332, PubMed: 9169480). Mediates

GM-CSF/CSF2-driven pain and inflammation (PubMed: <u>27525438</u>). In the brain, required to maintain the typical, highly branched morphology of hippocampal

microglia under homeostatic conditions. May be important for the

appropriate adaptation of microglial morphology and synaptic plasticity to acute lipopolysaccharide (LPS)-induced neuroinflammation (By similarity). Plays a role in wound healing, mainly by inducing fibroblast migration into the

wound (By similarity).

Cellular Location Secreted


Tissue Location Constitutively expressed in thymus. Detected at lower levels in the lung, colon

and small intestine (PubMed:8702936) Expressed in stimulated peripheral blood mononuclear cells, but not in resting cells (PubMed:8702936).

Background

Chemotactic factor for T-lymphocytes but not monocytes or granulocytes. May play a role in T-cell development in thymus and in trafficking and activation of mature T-cells. Binds to CCR4.

Images



Western blot analysis of various lysates, primary antibody was diluted at 1:1000, 4° over night, secondary antibody(cat: RS23920)was diluted at 1:10000, 37° 1hour.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.