

Polyclonal Antibody to Regulated On Activation In Normal T-Cell Expressed And Secreted (RANTES)

Regulated On Activation In Normal T-Cell Expressed And Secreted

Catalog # AP74507

Product Information

Application	WB
Primary Accession	P97272
Reactivity	Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	10088

Additional Information

Gene ID	100135495
Other Names	CCL5; SCYA5; SISd; EoCP; Chemokine C-C-Motif Ligand 5; Eosinophil Chemotactic Cytokine; T-cell Specific Protein p288; Small Inducible Cytokine Subfamily A(Cys-Cys)Member 5
Target/Specificity	Cavia (Guinea pig)
Dilution	WB~~Western blotting: 0.2-2 µg/mL;1:250-2500 Immunohistochemistry: 5-20 µg/mL;1:25-100 Immunocytochemistry: 5-20 µg/mL;1:25-100 Optimal working dilutions must be determined by end user.
Format	PBS, pH7.4, containing 0.02% NaN ₃ , 50% glycerol.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	CCL5
Synonyms	SCYA5
Function	Chemoattractant for blood monocytes, memory T-helper cells and eosinophils. Causes the release of histamine from basophils and activates eosinophils. May activate several chemokine receptors including CCR1, CCR3, CCR4 and CCR5. May also be an agonist of the G protein-coupled receptor GPR75. Together with GPR75, may play a role in neuron survival through activation of a downstream signaling pathway involving the PI3, Akt and MAP kinases. By activating GPR75 may also play a role in insulin secretion by islet

cells.

Cellular Location

Secreted.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.