

Polyclonal Antibody to Interleukin 11 (IL11)

Interleukin 11 Catalog # AP74535

Product Information


Application	WB
Primary Accession	<u>P47873</u>
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21522

Additional Information

Gene ID	16156
Other Names	AGIF; Adipogenesis Inhibitory Factor; Oprelvekin
Target/Specificity	Mus musculus (Mouse)
Dilution	WB~~Western blotting: 0.5-2 [g/mL;1:500-2000 Immunohistochemistry: 5-20 [g/mL;1:50-200 Immunocytochemistry: 5-20 [g/mL;1:50-200 Optimal working dilutions must be determined by end user.
Format	PBS, pH7.4, containing 0.02% NaN3, 50% glycerol.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	II11 {ECO:0000312 MGI:MGI:107613}
Function	Cytokine that stimulates the proliferation of hematopoietic stem cells and megakaryocyte progenitor cells and induces megakaryocyte maturation resulting in increased platelet production (PubMed: <u>8913282</u>). Also promotes the proliferation of hepatocytes in response to liver damage (PubMed: <u>22253262</u>). Binding to its receptor formed by IL6ST and either IL11RA1 or IL11RA2 activates a signaling cascade that promotes cell proliferation, also in the context of various cancers (PubMed: <u>10026196</u> , PubMed: <u>23948300</u>). Signaling leads to the activation of intracellular protein kinases and the phosphorylation of STAT3 (PubMed: <u>22253262</u> , PubMed: <u>23948300</u>). The interaction with the membrane- bound IL11RA and IL6ST stimulates 'classic signaling', whereas the binding of IL11 and soluble IL11RA to IL6ST stimulates 'trans- signaling' (By similarity).
Cellular Location	Secreted.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.