

Polyclonal Antibody to Interleukin 13 (IL13)

Interleukin 13

Catalog # AP74539

Product Information

Application	WB
Primary Accession	Q9N0W9
Reactivity	Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	14268

Additional Information

Gene ID	442990
Other Names	ALRH; BHR1; P600; IL-13; NC30
Target/Specificity	Canis familiaris; Canine (Dog)
Dilution	WB~~Western blotting: 0.5-2 μ g/mL; Immunohistochemistry: 5-20 μ g/mL; Immunocytochemistry: 5-20 μ g/mL; Optimal working dilutions must be determined by end user.
Format	PBS, pH7.4, containing 0.02% NaN ₃ , 50% glycerol.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	IL13
Function	Cytokine that plays important roles in allergic inflammation and immune response to parasite infection. Synergizes with IL2 in regulating interferon-gamma synthesis. Stimulates B-cell proliferation, and activation of eosinophils, basophils, and mast cells (By similarity). Plays an important role in controlling IL33 activity by modulating the production of transmembrane and soluble forms of interleukin-1 receptor-like 1/IL1RL1 (By similarity). Displays the capacity to antagonize Th1-driven proinflammatory immune response and downregulates synthesis of many proinflammatory cytokines including IL1, IL6, IL10, IL12 and TNF through a mechanism that partially involves suppression of NF-kappa-B (By similarity). Also functions on nonhematopoietic cells, including endothelial cells where it induces vascular cell adhesion protein 1/VCAM1, which is important in the recruitment of eosinophils. Exerts its biological effects through its receptors which comprises the IL4R chain and the IL13RA1 chain, to activate JAK1 and TYK2, leading to

the activation of STAT6. Aside from IL13RA1, another receptor IL13RA2 acts as a high affinity decoy for IL13 and mediates internalization and depletion of extracellular IL13 (By similarity). Induces inflammatory response in esophageal epithelium which results in a dysregulation of desmosome intercellular adhesion junctions, via down-regulation of DSG1 (By similarity).

Cellular Location

Secreted.

Images




Figure. Western Blot; Sample: Recombinant IL13, Canine.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.