

EHD3 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7458A

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	Q9NZN3
Other Accession	Q641Z6 , Q9WVK4 , Q9H4M9 , Q5E9R3
Reactivity	Human, Mouse
Predicted	Bovine, Mouse, Rat, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB17956
Calculated MW	60887
Antigen Region	22-48

Additional Information

Gene ID	30845
Other Names	EH domain-containing protein 3, PAST homolog 3, EHD3, EHD2, PAST3
Target/Specificity	This EHD3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 22-48 amino acids from the N-terminal region of human EHD3.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	EHD3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	EHD3 (HGNC:3244)
Function	ATP- and membrane-binding protein that controls membrane reorganization/tubulation upon ATP hydrolysis (PubMed: 25686250). In vitro

causes tubulation of endocytic membranes (PubMed:[24019528](#)). Binding to phosphatidic acid induces its membrane tubulation activity (By similarity). Plays a role in endocytic transport. Involved in early endosome to recycling endosome compartment (ERC), retrograde early endosome to Golgi, and endosome to plasma membrane (rapid recycling) protein transport. Involved in the regulation of Golgi maintenance and morphology (PubMed:[16251358](#), PubMed:[17233914](#), PubMed:[19139087](#), PubMed:[23781025](#)). Involved in the recycling of internalized D1 dopamine receptor (PubMed:[21791287](#)). Plays a role in cardiac protein trafficking probably implicating ANK2 (PubMed:[20489164](#)). Involved in the ventricular membrane targeting of SLC8A1 and CACNA1C and probably the atrial membrane localization of CACNA1GG and CACNA1H implicated in the regulation of atrial myocyte excitability and cardiac conduction (By similarity). In conjunction with EHD4 may be involved in endocytic trafficking of KDR/VEGFR2 implicated in control of glomerular function (By similarity). Involved in the rapid recycling of integrin beta-3 implicated in cell adhesion maintenance (PubMed:[23781025](#)). Involved in the unidirectional retrograde dendritic transport of endocytosed BACE1 and in efficient sorting of BACE1 to axons implicating a function in neuronal APP processing (By similarity). Plays a role in the formation of the ciliary vesicle, an early step in cilium biogenesis; possibly sharing redundant functions with EHD1 (PubMed:[25686250](#)).

Cellular Location

Recycling endosome membrane; Peripheral membrane protein; Cytoplasmic side. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, cilium membrane; Peripheral membrane protein; Cytoplasmic side. Note=Localizes to the ciliary pocket from where the cilium protrudes (PubMed:25686250) Colocalizes with RAB8A and MYO5B to a cytoplasmic tubular network devoid of RAB11A (By similarity). Colocalizes with ANK2 in myocyte perinuclear region (PubMed:20489164). Colocalizes with BACE1 in tubulovesicular cytoplasmic membranes. Colocalizes with BACE1 and APP amyloid beta proteins in hippocampal mossy fiber terminals (By similarity). {ECO:0000250|UniProtKB:Q9QXY6, ECO:0000269|PubMed:20489164, ECO:0000269|PubMed:25686250}

Tissue Location

Highly expressed in heart and brain and moderately expressed in kidney, liver, and placenta

Background



EHD3 plays a role in endocytic transport.

References


Soranzo, N., et al. Nat. Genet. 41(11):1182-1190(2009)
 Naslavsky, N., et al. J. Cell. Sci. 122 (PT 3), 389-400 (2009)
 Roland, J.T., et al. Mol. Biol. Cell 18(8):2828-2837(2007)

Images


Anti-EHD3 Antibody (N-term) at 1:1000 dilution + human brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 61 kDa
 Blocking/Dilution buffer: 5% NFDm/TBST.



Western blot analysis of EHD3 Antibody (N-term) (Cat. #AP7458a) in mouse lung tissue lysates (35ug/lane).EHD3 (arrow) was detected using the purified Pab.



EHD3 Antibody (N-term) (Cat. #AP7458a) IHC analysis in formalin fixed and paraffin embedded brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the EHD3 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



EHD3 Antibody (N-term) (Cat. #AP7458a) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.