

Polyclonal Antibody to Coagulation Factor VIII (F8)

Coagulation Factor VIII

Catalog # AP74659

Product Information

Application	WB
Primary Accession	O18806
Reactivity	Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	265831

Additional Information


Gene ID	403875
Other Names	FVIII; AHF; HEMA; F8B; F8C; Antihemophilic Globulin; Hemophilia A; Anti-Hemophilic Factor A; FVIII Procoagulant Component; F8 Protein
Target/Specificity	Canis familiaris; Canine (Dog)
Dilution	WB~~Western blotting: 0.5-2 μ g/mL Immunohistochemistry: 5-20 μ g/mL Immunocytochemistry: 5-20 μ g/mL Optimal working dilutions must be determined by end user.
Format	PBS, pH7.4, containing 0.02% NaN ₃ , 50% glycerol.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	F8
Function	Factor VIII, along with calcium and phospholipid, acts as a cofactor for factor IXa when it converts factor X to the activated form, factor Xa.
Cellular Location	Secreted, extracellular space.

Images

Figure. Western Blot; Sample: Recombinant F8, Canine.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.