

Polyclonal Antibody to Pleiotrophin (PTN)

Pleiotrophin

Catalog # AP74672

Product Information

Application	WB
Primary Accession	P63090
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	18869

Additional Information

Gene ID	24924
Other Names	HARP; HBGF8; HBNF; NEGF1; Heparin Binding Growth Factor 8; Neurite Growth-Promoting Factor 1; Heparin Affin Regulatory Peptide; Heparin Binding Growth Associated Molecule
Target/Specificity	Rattus norvegicus (Rat)
Dilution	WB~~Western blotting: 0.2-2 μ g/mL;1:250-2500 Immunohistochemistry: 5-20 μ g/mL;1:25-100 Immunocytochemistry: 5-20 μ g/mL;1:25-100 Optimal working dilutions must be determined by end user.
Format	PBS, pH7.4, containing 0.02% NaN ₃ , 50% glycerol.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	Ptn {ECO:0000312 RGD:3444}
Function	Secreted growth factor that mediates its signal through cell- surface proteoglycan and non-proteoglycan receptors (PubMed: 16814777 , PubMed: 27671118 , PubMed: 9817766). Binds cell-surface proteoglycan receptor via their chondroitin sulfate (CS) groups (By similarity). Thereby regulates many processes like cell proliferation, cell survival, cell growth, cell differentiation and cell migration in several tissues namely neuron and bone (PubMed: 27671118 , PubMed: 9817766). Also plays a role in synaptic plasticity and learning- related behavior by inhibiting long-term synaptic potentiation (By similarity). Binds PTPRZ1, leading to neutralization of the negative charges of the CS chains of PTPRZ1, inducing PTPRZ1 clustering, thereby causing the dimerization and inactivation of its phosphatase activity leading to increased

tyrosine phosphorylation of each of the PTPRZ1 substrates like ALK or AFAP1L2 in order to activate the PI3K-AKT pathway (PubMed:[16814777](#)). Through PTPRZ1 binding, regulates endothelial cell migration and controls oligodendrocyte precursor cell differentiation (By similarity). Forms a complex with PTPRZ1 and integrin alpha-V/beta-3 (ITGAV:ITGB3) that stimulates endothelial cell migration through SRC dephosphorylation and activation that consequently leads to ITGB3 'Tyr-773' phosphorylation (By similarity). In adult hippocampus promotes dendritic arborization, spine development, and functional integration and connectivity of newborn granule neurons through ALK by activating AKT signaling pathway (By similarity). Binds GPC2 and chondroitin sulfate proteoglycans (CSPGs) at the neuron surface, leading to abrogation of binding between PTPRS and CSPGs and neurite outgrowth promotion (PubMed:[27671118](#)). Binds SDC3 and mediates bone formation by recruiting and attaching osteoblasts/osteoblast precursors to the sites for new bone deposition (PubMed:[9817766](#)). Binds ALK and promotes cell survival and cell proliferation through MAPK pathway activation (By similarity). Inhibits proliferation and enhances differentiation of neural stem cells by inhibiting FGF2-induced fibroblast growth factor receptor signaling pathway (By similarity). Mediates regulatory mechanisms in normal hemostasis and in hematopoietic regeneration and in maintaining the balance of myeloid and lymphoid regeneration (By similarity). In addition may play a role in the female reproductive system, auditory response and the progesterone-induced decidualization pathway (By similarity).


Cellular Location

Secreted {ECO:0000250|UniProtKB:P21246}.

Tissue Location

Expressed in brain (PubMed:2170351). Hardly expressed in bone. Abundantly expressed in the growth plate. Intensely expressed in the cartilage matrix that forms the hollow for the secondary ossification center. Is expressed selectively by the osteocytes of the lamellar bone, whereas it is hardly expressed by those of the woven bone. Is detected in the matrix surrounding the osteocytes and in the canaliculi of the osteocytes. Is distributed on the surface of lamellar bone (PubMed:9817766)

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.