



Polyclonal Antibody to Chemokine (C-X-C Motif) Ligand 3 (CXCL3)

Chemokine (C-X-C Motif) Ligand 3 Catalog # AP74701

Product Information

ApplicationWBPrimary AccessionQ10746ReactivityRatHostRabbitClonalityPolyclonalCalculated MW11109

Additional Information

Gene ID 171551

Other Names GROg; SCYB3; GRO3; GRO-G; MIP2-B; MIP2b; CINC2b; CINC2b; Growth

Regulated Oncogene Gamma; Macrophage inflammatory protein 2-beta

Target/Specificity Rattus norvegicus (Rat)

Dilution WB~~Western blotting: 0.2-2 \(\text{Ig/mL}; 1:250-2500 \)

Immunohistochemistry: 5-20 [g/mL;1:25-100 Immunocytochemistry: 5-20 [g/mL;1:25-100

Optimal working dilutions must be determined by end user.

Format PBS, pH7.4, containing 0.02% NaN3, 50% glycerol.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name Cxcl3

Synonyms Cinc2


Function Ligand for CXCR2 (By similarity). Has chemotactic activity for neutrophils.

May play a role in inflammation and exert its effects on endothelial cells in an

autocrine fashion.

Cellular Location Secreted.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.