

Polyclonal Antibody to Transition Protein 2 (TNP2)

Transition Protein 2

Catalog # AP74767

Product Information

Application	WB
Primary Accession	P11378
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	13117

Additional Information

Gene ID	21959
Other Names	TP2; During Histone To Protamine Replacement; Nuclear transition protein 2
Target/Specificity	Mus musculus (Mouse)
Dilution	WB~~Western blotting: 0.5-2 μ g/mL; Immunohistochemistry: 5-20 μ g/mL; Immunocytochemistry: 5-20 μ g/mL; Optimal working dilutions must be determined by end user.
Format	PBS, pH7.4, containing 0.02% NaN ₃ , 50% glycerol.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	Tnp2 {ECO:0000312 MGI:MGI:98785}
Function	Plays a key role in the replacement of histones to protamine in the elongating spermatids of mammals (PubMed: 15083521 , PubMed: 15163613 , PubMed: 15189834 , PubMed: 28366643). In condensing spermatids, loaded onto the nucleosomes, where it promotes the recruitment and processing of protamines, which are responsible for histone eviction (PubMed: 28366643). The histone H2AB1-H2BC1/TH2B dimer is required for loading of TNP2 onto chromatin (PubMed: 28366643).
Cellular Location	Nucleus. Chromosome. Note=Loaded onto the nucleosomes of condensing spermatids (PubMed:28366643). Inclusion of the H2AB1- H2BC1/TH2B dimer into chromatin opens the nucleosomes, releasing the nucleosomal DNA ends and allowing the invasion of nucleosomes by transition protein TNP2 (PubMed:28366643). Nuclear import is mediated by IPO4. Nucleolar localization requires the protein to be phosphorylated (By similarity).

Images




Figure. Western Blot; Sample: Recombinant TNP2, Mouse.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.