

# Thioredoxin Rabbit mAb

Catalog # AP74806

## **Product Information**

**Application** WB, IHC-P, IHC-F, IP, ICC

Primary Accession P10599
Reactivity Human
Host Rabbit

**Clonality** Monoclonal Antibody

Calculated MW 11737

### **Additional Information**

**Gene ID** 7295

Other Names TXN

**Dilution** WB~~1/1000-1/10000 IHC-P~~N/A IHC-F~~N/A IP~~N/A ICC~~N/A

Format Liquid

#### **Protein Information**

Name TXN

Synonyms TRDX, TRX, TRX1

**Function** Participates in various redox reactions through the reversible oxidation of its

active center dithiol to a disulfide and catalyzes dithiol-disulfide exchange reactions (PubMed:17182577, PubMed:19032234, PubMed:2176490). Plays a role in the reversible S- nitrosylation of cysteine residues in target proteins, and thereby contributes to the response to intracellular nitric oxide. Nitrosylates the active site Cys of CASP3 in response to nitric oxide (NO), and

thereby inhibits caspase-3 activity (PubMed:<u>16408020</u>, PubMed:<u>17606900</u>). Induces the FOS/JUN AP-1 DNA-binding activity in ionizing radiation (IR) cells through its oxidation/reduction status and stimulates AP-1 transcriptional

activity (PubMed: 11118054, PubMed: 9108029).


**Cellular Location**Nucleus. Cytoplasm. Secreted Note=Translocates from the cytoplasm into the


nucleus after phorbol 12- myristate 13-acetate induction (PMA)


(PubMed:9108029). Predominantly in the cytoplasm in non irradiated cells (PubMed:11118054). Radiation induces translocation of TRX from the cytoplasm to the nucleus (PubMed:11118054). Secreted by a leaderless

secretory pathway (PubMed:1332947).

# **Images**







Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.