

MCM6 Rabbit mAb

Catalog # AP74817

Product Information

Application	WB, IHC-P, IHC-F, IP, ICC
Primary Accession	Q14566
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	92889




Additional Information

Gene ID	4175
Other Names	MCM6
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IP~~N/A ICC~~N/A
Format	Liquid

Protein Information

Name	MCM6 (HGNC:6949)
Function	Acts as a component of the MCM2-7 complex (MCM complex) which is the replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. Core component of CDC45-MCM-GINS (CMG) helicase, the molecular machine that unwinds template DNA during replication, and around which the replisome is built (PubMed: 16899510 , PubMed: 32453425 , PubMed: 34694004 , PubMed: 34700328 , PubMed: 35585232 , PubMed: 9305914). The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity (PubMed: 32453425).
Cellular Location	Nucleus. Chromosome. Note=Binds to chromatin during G1 and detaches from it during S phase.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.