

[KO]FGF2 Rabbit mAb

Catalog # AP74819

Product Information

Application	WB, FC, IP
Primary Accession	P09038
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	30770

Additional Information

Gene ID	2247
Other Names	FGF2
Dilution	WB~~1:500-1:1000 FC~~1:200-1:500 IP~~1:20-1:50
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	FGF2
Synonyms	FGFB
Function	Acts as a ligand for FGFR1, FGFR2, FGFR3 and FGFR4 (PubMed: 8663044). Also acts as an integrin ligand which is required for FGF2 signaling (PubMed: 28302677). Binds to integrin ITGAV:ITGB3 (PubMed: 28302677). Plays an important role in the regulation of cell survival, cell division, cell differentiation and cell migration (PubMed: 28302677 , PubMed: 8663044). Functions as a potent mitogen in vitro (PubMed: 1721615 , PubMed: 3732516 , PubMed: 3964259). Can induce angiogenesis (PubMed: 23469107 , PubMed: 28302677). Mediates phosphorylation of ERK1/2 and thereby promotes retinal lens fiber differentiation (PubMed: 29501879).
Cellular Location	Secreted. Nucleus. Note=Exported from cells by an endoplasmic reticulum (ER)/Golgi-independent mechanism. Unconventional secretion of FGF2 occurs by direct translocation across the plasma membrane (PubMed:20230531).

Binding of exogenous FGF2 to FGFR facilitates endocytosis followed by translocation of FGF2 across endosomal membrane into the cytosol (PubMed:22321063). Nuclear import from the cytosol requires the classical nuclear import machinery, involving proteins KPNA1 and KPNB1, as well as CEP57 (PubMed:22321063)

Tissue Location


Expressed in granulosa and cumulus cells. Expressed in hepatocellular carcinoma cells, but not in non-cancerous liver tissue.

Background


FGF2 is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. FGF2 is a single-chain polypeptide growth factor that plays a significant role in the process of wound healing and is a potent inducer of angiogenesis. Due to its basic pH, the factor is named FGF-2 (basic FGF, bFGF). Several different forms of the human protein exist ranging from 18-24 kDa in size due to the use of alternative start sites within the fgf-2 gene. It has a 55 percent amino acid residue identity to FIBROBLAST GROWTH FACTOR 1 and has potent heparin-binding activity. The growth factor is an extremely potent inducer of DNA synthesis in a variety of cell types from mesoderm and neuroectoderm lineages. It was originally named basic fibroblast growth factor based upon its chemical properties and to distinguish it from acidic fibroblast growth factor (FIBROBLAST GROWTH FACTOR 1).



Images

Western blot analysis of FGF2 in HeLa(WT) and FGF2-Knockout HeLa(KO) using FGF2 antibody.



Western blot analysis of FGF2 in HeLa lysates using FGF2 antibody.





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.