

# IDH1 Rabbit mAb

Catalog # AP74833

## Product Information

---

<b>Application</b>	WB, IHC-P, IP
<b>Primary Accession</b>	<a href="#">O75874</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Calculated MW</b>	46659

## Additional Information

---

<b>Gene ID</b>	3417
<b>Other Names</b>	IDH1
<b>Dilution</b>	WB~~1/500-1/1000 IHC-P~~N/A IP~~N/A
<b>Format</b>	Liquid



## Protein Information

---

<b>Name</b>	IDH1
<b>Synonyms</b>	PICD
<b>Function</b>	Catalyzes the NADP(+)-dependent oxidative decarboxylation of isocitrate (D-threo-isocitrate) to 2-ketoglutarate (2-oxoglutarate), which is required by other enzymes such as the phytanoyl-CoA dioxygenase (PubMed: <a href="#">10521434</a> , PubMed: <a href="#">19935646</a> ). Plays a critical role in the generation of NADPH, an important cofactor in many biosynthesis pathways (PubMed: <a href="#">10521434</a> ). May act as a corneal epithelial crystallin and may be involved in maintaining corneal epithelial transparency (By similarity).
<b>Cellular Location</b>	Cytoplasm, cytosol. Peroxisome

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.