

TAF3 Rabbit mAb

Catalog # AP74840

Product Information

Application WB, IHC-P
Primary Accession Q5VWG9
Reactivity Human
Rabbit

Clonality Monoclonal Antibody

Calculated MW 103582

Additional Information

Gene ID 83860

Other Names TAF3

Dilution WB~~1/500-1/1000 IHC-P~~N/A

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Protein Information

Name TAF3

Function The TFIID basal transcription factor complex plays a major role in the

initiation of RNA polymerase II (Pol II)-dependent transcription


(PubMed:33795473). TFIID recognizes and binds promoters with or without a TATA box via its subunit TBP, a TATA-box-binding protein, and promotes assembly of the pre-initiation complex (PIC) (PubMed:33795473). The TFIID complex consists of TBP and TBP-associated factors (TAFs), including TAF1, TAF2, TAF3, TAF4, TAF5, TAF6, TAF7, TAF8, TAF9, TAF10, TAF11, TAF12 and TAF13 (PubMed:33795473). The TFIID complex structure can be divided into 3 modules TFIID-A, TFIID-B, and TFIID-C (PubMed:33795473). TAF3 forms the TFIID-A module together with TAF5 and TBP (PubMed:33795473). Required in complex with TBPL2 for the differentiation of myoblasts into myocytes (PubMed:11438666). The TAF3-TBPL2 complex replaces TFIID at specific


promoters at an early stage in the differentiation process

(PubMed: 11438666).

Cellular Location Nucleus.

Images





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.