

[KD]CENPA Rabbit mAb

Catalog # AP74853

Product Information

Application	WB, IHC-P
Primary Accession	P49450
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	15991

Additional Information

Gene ID	1058
Other Names	CENPA
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	CENPA
Function	Histone H3-like nucleosomal protein that is specifically found in centromeric nucleosomes (PubMed:11756469, PubMed:14667408, PubMed:15282608, PubMed:15475964, PubMed:15702419, PubMed:17651496, PubMed:19114591, PubMed:20739937, PubMed:27499292, PubMed:7962047, PubMed:9024683). Replaces conventional H3 in the nucleosome core of centromeric chromatin that serves as an assembly site for the inner kinetochore (PubMed:18072184). The presence of CENPA subtly modifies the nucleosome structure and the way DNA is wrapped around the nucleosome and gives rise to protruding DNA ends that are less well- ordered and rigid compared to nucleosomes containing histone H3 (PubMed:26878239, PubMed:27499292). May serve as an epigenetic mark that propagates centromere identity through replication and cell division (PubMed:15282608, PubMed:15475964, PubMed:20739937, PubMed:21478274, PubMed:26878239). Required for recruitment and assembly of kinetochore proteins, and as a consequence required for progress through mitosis,

chromosome segregation and cytokinesis (PubMed:[11756469](#), PubMed:[14667408](#), PubMed:[18072184](#), PubMed:[23818633](#), PubMed:[25556658](#), PubMed:[27499292](#)).


Cellular Location

Nucleus. Chromosome, centromere. Note=Localizes exclusively to sites of kinetochore assembly in centromeres. Occupies a compact domain at the inner kinetochore plate stretching across 2 thirds of the length of the constriction but encompassing only one third of the constriction width and height (PubMed:19114591) Phosphorylation at Ser-68 during early mitosis abolishes association with chromatin and centromeres and results in dispersed nuclear location (PubMed:25556658).

Background

Centromeres are the differentiated chromosomal domains that specify the mitotic behavior of chromosomes. This gene encodes a centromere protein which contains a histone H3 related histone fold domain that is required for targeting to the centromere. Centromere protein A is proposed to be a component of a modified nucleosome or nucleosome-like structure in which it replaces 1 or both copies of conventional histone H3 in the (H3-H4)₂ tetrameric core of the nucleosome particle. The protein is a replication-independent histone that is a member of the histone H3 family. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Images



Western blot analysis of CENPA in K562, HeLa lysates using CENPA antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.