10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

FDFT1 Rabbit mAb

Catalog # AP74900

Product Information

Application	WB, IHC-P, IP
Primary Accession	P37268
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	48115

Additional Information

Gene ID	2222
Other Names	FDFT1
Dilution	WB~~1:500-1:1000 IHC-P~~N/A IP~~1:50-1:100
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.


Protein Information

Name	FDFT1
Function	Catalyzes the condensation of 2 farnesyl pyrophosphate (FPP) moieties to form squalene. Proceeds in two distinct steps. In the first half-reaction, two molecules of FPP react to form the stable presqualene diphosphate intermediate (PSQPP), with concomitant release of a proton and a molecule of inorganic diphosphate. In the second half-reaction, PSQPP undergoes heterolysis, isomerization, and reduction with NADPH or NADH to form squalene. It is the first committed enzyme of the sterol biosynthesis pathway.
Cellular Location	Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q02769}; Multi-pass membrane protein
Tissue Location	Widely expressed..


Background

Critical branch point enzyme of isoprenoid biosynthesis that is thought to regulate the flux of isoprene intermediates through the sterol pathway.


Images




Western blot analysis of FDFT1 in K562, C6, Hela lysates using FDFT1 antibody.



Immunocytochemistry analysis of FDFT1(green) in Hela using FDFT1 antibody,and DAPI(blue)





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.