

# PMS2 Rabbit mAb

Catalog # AP74907

## Product Information

---

Application	WB, IP
Primary Accession	<a href="#">P54278</a>
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	95797

## Additional Information

---

Gene ID	5395
Other Names	PMS2
Dilution	WB~~1/500-1/1000 IP~~N/A
Format	Liquid


## Protein Information

---

Name	PMS2 ( <a href="#">HGNC:9122</a> )
Function	<p>Component of the post-replicative DNA mismatch repair system (MMR) (PubMed:<a href="#">30653781</a>, PubMed:<a href="#">35189042</a>). Heterodimerizes with MLH1 to form MutL alpha. DNA repair is initiated by MutS alpha (MSH2-MSH6) or MutS beta (MSH2-MSH3) binding to a dsDNA mismatch, then MutL alpha is recruited to the heteroduplex. Assembly of the MutL-MutS-heteroduplex ternary complex in presence of RFC and PCNA is sufficient to activate endonuclease activity of PMS2. It introduces single-strand breaks near the mismatch and thus generates new entry points for the exonuclease EXO1 to degrade the strand containing the mismatch. DNA methylation would prevent cleavage and therefore assure that only the newly mutated DNA strand is going to be corrected. MutL alpha (MLH1-PMS2) interacts physically with the clamp loader subunits of DNA polymerase III, suggesting that it may play a role to recruit the DNA polymerase III to the site of the MMR. Also implicated in DNA damage signaling, a process which induces cell cycle arrest and can lead to apoptosis in case of major DNA damages. Possesses an ATPase activity, but in the absence of gross structural changes, ATP hydrolysis may not be necessary for proficient mismatch repair (PubMed:<a href="#">35189042</a>).</p>
Cellular Location	Nucleus

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.