

Glutamine Synthetase Rabbit mAb

Catalog # AP74913

Product Information

ApplicationWBPrimary AccessionP15104ReactivityMouse, RatHostRabbit

Clonality Monoclonal Antibody

Calculated MW 42064

Additional Information

Gene ID 2752

Other Names GLUL

Dilution WB~~1/500-1/1000

Format Liquid

Protein Information

Name GLUL {ECO:0000303 | PubMed:30158707, ECO:0000312 | HGNC:4341}

Function Glutamine synthetase that catalyzes the ATP-dependent conversion of

glutamate and ammonia to glutamine (PubMed: 16267323, PubMed: 30158707, PubMed: 36289327). Its role depends on tissue

localization: in the brain, it regulates the levels of toxic ammonia and converts neurotoxic glutamate to harmless glutamine, whereas in the liver, it is one of the enzymes responsible for the removal of ammonia (By similarity). Plays a key role in ammonium detoxification during erythropoiesis: the glutamine synthetase activity is required to remove ammonium generated by porphobilinogen deaminase (HMBS) during heme biosynthesis to prevent ammonium accumulation and oxidative stress (By similarity). Essential for proliferation of fetal skin fibroblasts (PubMed: 18662667). Independently of its glutamine synthetase activity, required for endothelial cell migration during vascular development; acts by regulating membrane localization and activation of the GTPase RHOJ, possibly by promoting RHOJ palmitoylation (PubMed:30158707). May act as a palmitoyltransferase for RHOJ: able to autopalmitoylate and then transfer the palmitoyl group to RHOJ (PubMed:30158707). Plays a role in ribosomal 40S subunit biogenesis (PubMed: <u>26711351</u>). Through the interaction with BEST2, inhibits BEST2 channel activity by affecting the gating at the aperture in the absence of intracellular L-glutamate, but sensitizes BEST2 to intracellular L-glutamate, which promotes the opening of BEST2 and thus relieves its inhibitory effect on BEST2 (PubMed: 36289327).

Cytoplasm, cytosol. Microsome {ECO:0000250 | UniProtKB:P09606}

Mitochondrion {ECO:0000250 | UniProtKB:P09606}. Cell membrane;


Lipid-anchor. Note=Mainly localizes in the cytosol, with a fraction associated


with the cell membrane

Tissue Location

Expressed in endothelial cells.

Images





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.