

ATG7 Rabbit mAb

Catalog # AP75010

Product Information

Application	WB, IP
Primary Accession	O95352
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	77960

Additional Information

Gene ID	10533
Other Names	ATG7
Dilution	WB~~1/500-1/1000 IP~~N/A
Format	Liquid

Protein Information


Name	ATG7 (HGNC:16935)
Synonyms	APG7L
Function	<p>E1-like activating enzyme involved in the 2 ubiquitin-like systems required for cytoplasm to vacuole transport (Cvt) and autophagy. Activates ATG12 for its conjugation with ATG5 as well as the ATG8 family proteins for their conjugation with phosphatidylethanolamine. Both systems are needed for the ATG8 association to Cvt vesicles and autophagosomes membranes. Required for autophagic death induced by caspase-8 inhibition. Facilitates LC3-I lipidation with phosphatidylethanolamine to form LC3-II which is found on autophagosomal membranes (PubMed:34161705). Required for mitophagy which contributes to regulate mitochondrial quantity and quality by eliminating the mitochondria to a basal level to fulfill cellular energy requirements and preventing excess ROS production. Modulates p53/TP53 activity to regulate cell cycle and survival during metabolic stress. Also plays a key role in the maintenance of axonal homeostasis, the prevention of axonal degeneration, the maintenance of hematopoietic stem cells, the formation of Paneth cell granules, as well as in adipose differentiation. Plays a role in regulating the liver clock and glucose metabolism by mediating the autophagic degradation of CRY1 (clock repressor) in a time-dependent manner (By similarity).</p>
Cellular Location	Cytoplasm. Preautophagosomal structure. Note=Also localizes to discrete

punctae along the ciliary axoneme and to the base of the ciliary axoneme

Tissue Location

Widely expressed, especially in kidney, liver, lymph nodes and bone marrow.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.