

ACADM Rabbit mAb

Catalog # AP75023

Product Information

Primary Accession	P11310
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	46588




Additional Information

Gene ID	34
Other Names	ACADM
Format	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	ACADM (HGNC:89)
Function	Medium-chain specific acyl-CoA dehydrogenase is one of the acyl-CoA dehydrogenases that catalyze the first step of mitochondrial fatty acid beta-oxidation (FAO), breaking down fatty acids into acetyl- CoA and allowing the production of energy from fats (PubMed:1970566, PubMed:21237683, PubMed:2251268, PubMed:8823175). The first step of FAO consists in the proR-proR stereospecific alpha, beta-dehydrogenation of fatty acyl-CoA thioesters using the electron transfer flavoprotein (ETF) as their physiologic electron acceptor, resulting in the formation of trans-2-enoyl-CoA ((2E)-enoyl-CoA) (PubMed:2251268). ETF is the electron acceptor that transfers electrons to the main mitochondrial respiratory chain via ETF-ubiquinone oxidoreductase (ETF dehydrogenase) (PubMed:15159392, PubMed:25416781). Among the different mitochondrial acyl-CoA dehydrogenases, medium-chain specific acyl-CoA dehydrogenase has preference for fatty acyl-CoAs with saturated 6 to 12 carbons long primary chains, making it but can also catalyze longer chains such as C14 and C16 (PubMed:1970566, PubMed:21237683, PubMed:2251268, PubMed:8823175).
Cellular Location	Mitochondrion matrix
Tissue Location	Expressed ubiquitously with highest levels in heart and muscle.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.