

# ACAT1 Rabbit mAb

Catalog # AP75025

## Product Information

---

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IP                 |
| Primary Accession | <a href="#">P24752</a> |
| Reactivity        | Human                  |
| Host              | Rabbit                 |
| Clonality         | Monoclonal Antibody    |
| Calculated MW     | 45200                  |

## Additional Information

---

|             |   |
|-------------|---|
| Gene ID     | 38  |
| Other Names | ACAT1   |
| Dilution    | WB~~1/500-1/1000 IP~~N/A  |
| Format      | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. |


## Protein Information

---

|                   |  |
|-------------------|--|
| Name              | ACAT1  |
| Synonyms          | ACAT, MAT  |
| Function          | <p>This is one of the enzymes that catalyzes the last step of the mitochondrial beta-oxidation pathway, an aerobic process breaking down fatty acids into acetyl-CoA (PubMed:<a href="#">1715688</a>, PubMed:<a href="#">7728148</a>, PubMed:<a href="#">9744475</a>). Using free coenzyme A/CoA, catalyzes the thiolytic cleavage of medium- to long-chain 3-oxoacyl-CoAs into acetyl-CoA and a fatty acyl-CoA shortened by two carbon atoms (PubMed:<a href="#">1715688</a>, PubMed:<a href="#">7728148</a>, PubMed:<a href="#">9744475</a>). The activity of the enzyme is reversible and it can also catalyze the condensation of two acetyl-CoA molecules into acetoacetyl-CoA (PubMed:<a href="#">17371050</a>). Thereby, it plays a major role in ketone body metabolism (PubMed:<a href="#">1715688</a>, PubMed:<a href="#">17371050</a>, PubMed:<a href="#">7728148</a>, PubMed:<a href="#">9744475</a>).</p> |
| Cellular Location | Mitochondrion.   |

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.