

CaMKII alpha Rabbit mAb

Catalog # AP75183

Product Information

Application	WB, IHC-P, FC
Primary Accession	Q9UQM7
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	54088

Additional Information

Gene ID	815
Other Names	CAMK2A
Dilution	WB~~1:1000-1:2000 IHC-P~~N/A FC~~1:20-1:100
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	CAMK2A
Synonyms	CAMKA, KIAA0968
Function	Calcium/calmodulin-dependent protein kinase that functions autonomously after Ca(2+)/calmodulin-binding and autophosphorylation, and is involved in various processes, such as synaptic plasticity, neurotransmitter release and long-term potentiation (PubMed: 14722083). Member of the NMDAR signaling complex in excitatory synapses, it regulates NMDAR-dependent potentiation of the AMPAR and therefore excitatory synaptic transmission (By similarity). Regulates dendritic spine development (PubMed: 28130356). Also regulates the migration of developing neurons (PubMed: 29100089). Phosphorylates the transcription factor FOXO3 to activate its transcriptional activity (PubMed: 23805378). Phosphorylates the transcription factor ETS1 in response to calcium signaling, thereby decreasing ETS1 affinity for DNA (By similarity). In response to interferon-gamma (IFN-gamma) stimulation, catalyzes phosphorylation of STAT1, stimulating the JAK- STAT signaling pathway

(PubMed:[11972023](#)). In response to interferon- beta (IFN-beta) stimulation, stimulates the JAK-STAT signaling pathway (PubMed:[35568036](#)). In response to interferon-gamma (IFN-gamma) stimulation, catalyzes phosphorylation of PSAT1, inhibiting ferroptosis by promoting GPX4 hydroxylation and stability (PubMed:[40281343](#)). Acts as a negative regulator of 2-arachidonoylglycerol (2-AG)-mediated synaptic signaling via modulation of DAGLA activity (By similarity).


Cellular Location

Synapse {ECO:0000250|UniProtKB:P11275}. Postsynaptic density {ECO:0000250|UniProtKB:P11275}. Cell projection, dendritic spine. Cell projection, dendrite. Note=Postsynaptic lipid rafts {ECO:0000250|UniProtKB:P11275}


Background

CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity.


Images



Western blot analysis of CaMKII alpha in 3T3 lysates using CaMKII alpha antibody.



Western blot analysis of CaMKII alpha in rat Brain, HeLa lysates using CaMKII alpha antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.