10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CD11a Rabbit mAb

Catalog # AP75209

Product Information

Application	WB, IHC-P, IHC-F, FC, IP
Primary Accession	P20701
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	128770

Additional Information

Gene ID	3683
Other Names	ITGAL
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A IHC-F~~N/A FC~~1:20-1:50 IP~~1:20-1:50
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	ITGAL (HGNC:6148)
Function	Integrin ITGAL/ITGB2 is a receptor for ICAM1, ICAM2, ICAM3 and ICAM4 (PubMed:10846180, PubMed:12526797, PubMed:1676048, PubMed:3086451). Integrin ITGAL/ITGB2 is a receptor for F11R (PubMed:11812992, PubMed:15528364). Integrin ITGAL/ITGB2 is a receptor for the secreted form of ubiquitin-like protein ISG15; the interaction is mediated by ITGAL (PubMed:29100055). Involved in a variety of immune phenomena including leukocyte-endothelial cell interaction, cytotoxic T-cell mediated killing, and antibody dependent killing by granulocytes and monocytes. Contributes to natural killer cell cytotoxicity (PubMed:15356110). Involved in leukocyte adhesion and transmigration of leukocytes including T-cells and neutrophils (PubMed:11812992). Acts as a platform at the immunological synapse to translate TCR engagement and density of the ITGAL ligand ICAM1 into graded adhesion (PubMed:38195629). Required for generation of common lymphoid progenitor cells in bone marrow, indicating a role in lymphopoiesis (By similarity). Integrin ITGAL/ITGB2 in association with ICAM3, contributes to

	apoptotic neutrophil phagocytosis by macrophages (PubMed:23775590). Integrins ITGAL:ITGB2 functions as a receptor of the neuron-specific adhesion molecule ICAM5 ensuring neuron cell-leukocyte adhesion (PubMed:10741396). Integrin ITGAL/ITGB2 that functions as a signaling receptor of ICAM2, ensuring leukocyte cell-cell adhesion on resting cells (PubMed:1676048).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Membrane raft. Note=Upon antigen recognition by the TCR, is recruited to lipid rafts (PubMed:15684041).
Tissue Location	Leukocytes.


Background


CD11a is a 170-180 kD type I transmembrane glycoprotein also known as LFA-1α chain and integrin αL subunit. CD11a non-covalently associates with integrin β2 (CD18) to form LFA-1. It is expressed on all leukocytes, including B and T lymphocytes, monocytes, macrophages, neutrophils, basophils and eosinophils. It is absent on non-hematopoietic tissues and platelets. CD11a plays a central role in leukocyte cell-cell interactions and is important in lymphocyte costimulation. CD11a/CD18 binds to ICAM-1 (CD54), ICAM-2 (CD102), and ICAM-3 (CD50).


Images



Western blot analysis of CD11a in THP-1 lysates using Integrin alpha L antibody.







Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.