

CD166 Rabbit mAb

Catalog # AP75212

Product Information

| | |
|--------------------------|--------------------------|
| Application | WB, IHC-P, IHC-F, FC, IP |
| Primary Accession | Q13740 |
| Reactivity | Rat, Human, Mouse |
| Host | Rabbit |
| Clonality | Monoclonal Antibody |
| Isotype | IgG |
| Conjugate | Unconjugated |
| Purification | Affinity Purified |
| Calculated MW | 65102 |

Additional Information

| | |
|--------------------|---|
| Gene ID | 214 |
| Other Names | ALCAM |
| Dilution | WB~~1:1000-1:5000 IHC-P~~N/A IHC-F~~N/A FC~~1:50-1:100 IP~~1:20-1:50 |
| Format | Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |

Protein Information

| | |
|-----------------|--|
| Name | ALCAM |
| Synonyms | MEMD {ECO:0000303 PubMed:9502422} |
| Function | Cell adhesion molecule that mediates both heterotypic cell- cell contacts via its interaction with CD6, as well as homotypic cell- cell contacts (PubMed: 15048703 , PubMed: 15496415 , PubMed: 16352806 , PubMed: 23169771 , PubMed: 24945728 , PubMed: 7760007). Promotes T-cell activation and proliferation via its interactions with CD6 (PubMed: 15048703 , PubMed: 16352806 , PubMed: 24945728). Contributes to the formation and maturation of the immunological synapse via its interactions with CD6 (PubMed: 15294938 , PubMed: 16352806). Mediates homotypic interactions with cells that express ALCAM (PubMed: 15496415 , PubMed: 16352806). Acts as a ligand for the LILRB4 receptor, enhancing LILRB4-mediated inhibition of T cell proliferation (PubMed: 29263213). Required for normal hematopoietic stem cell engraftment in the bone marrow (PubMed: 24740813). Mediates attachment of dendritic cells onto endothelial cells via homotypic interaction |

(PubMed:[23169771](#)). Inhibits endothelial cell migration and promotes endothelial tube formation via homotypic interactions (PubMed:[15496415](#), PubMed:[23169771](#)). Required for normal organization of the lymph vessel network. Required for normal hematopoietic stem cell engraftment in the bone marrow. Plays a role in hematopoiesis; required for normal numbers of hematopoietic stem cells in bone marrow. Promotes in vitro osteoblast proliferation and differentiation (By similarity). Promotes neurite extension, axon growth and axon guidance; axons grow preferentially on surfaces that contain ALCAM. Mediates outgrowth and pathfinding for retinal ganglion cell axons (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell projection, axon {ECO:0000250|UniProtKB:Q61490}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q61490}. Note=Detected at the immunological synapse, i.e, at the contact zone between antigen-presenting dendritic cells and T-cells (PubMed:15294938, PubMed:16352806). Colocalizes with CD6 and the TCR/CD3 complex at the immunological synapse (PubMed:15294938).


Tissue Location

Detected on hematopoietic stem cells derived from umbilical cord blood (PubMed:24740813). Detected on lymph vessel endothelial cells, skin and tonsil (PubMed:23169771). Detected on peripheral blood monocytes (PubMed:15048703). Detected on monocyte- derived dendritic cells (at protein level) (PubMed:16352806). Detected at low levels in spleen, placenta, liver (PubMed:9502422). Expressed by activated T-cells, B-cells, monocytes and thymic epithelial cells (PubMed:7760007). Isoform 1 and isoform 3 are detected in vein and artery endothelial cells, astrocytes, keratinocytes and artery smooth muscle cells (PubMed:15496415). Expressed by neurons in the brain Restricted expression in tumor cell lines. Detected in highly metastasizing melanoma cell lines (PubMed:9502422)

Background


Cell adhesion molecule that binds to CD6. Involved in neurite extension by neurons via heterophilic and homophilic interactions. May play a role in the binding of T- and B-cells to activated leukocytes, as well as in interactions between cells of the nervous system.

Images




Immunocytochemistry analysis of CD166 (green) in SKOV-3 using CD166 antibody, and DAPI(blue).



Western blot analysis of CD166 in mouse brain lysates using CD166 antibody.



Western blot analysis of CD166 in rat Brain, C6 lysates using CD166 antibody



Immunohistochemistry analysis of paraffin-embedded Human breast cancer using CD166 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.