

# Chk1 Rabbit mAb

Catalog # AP75258

## Product Information

---

|                          |                        |
|--------------------------|------------------------|
| <b>Application</b>       | WB, FC, IP             |
| <b>Primary Accession</b> | <a href="#">O14757</a> |
| <b>Reactivity</b>        | Rat, Human, Mouse      |
| <b>Host</b>              | Rabbit                 |
| <b>Clonality</b>         | Monoclonal Antibody    |
| <b>Isotype</b>           | IgG                    |
| <b>Conjugate</b>         | Unconjugated           |
| <b>Purification</b>      | Affinity Purified      |
| <b>Calculated MW</b>     | 54434                  |

## Additional Information

---

|                    |   |
|--------------------|---|
| <b>Gene ID</b>     | 1111  |
| <b>Other Names</b> | CHEK1   |
| <b>Dilution</b>    | WB~~1:500-1:1000 FC~~1:20 IP~~1:20  |
| <b>Format</b>      | Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. |
| <b>Storage</b>     | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.        |

## Protein Information

---

|                 |  |
|-----------------|--|
| <b>Name</b>     | CHEK1  |
| <b>Synonyms</b> | CHK1   |
| <b>Function</b> | Serine/threonine-protein kinase which is required for checkpoint-mediated cell cycle arrest and activation of DNA repair in response to the presence of DNA damage or unreplicated DNA (PubMed: <a href="#">11535615</a> , PubMed: <a href="#">12399544</a> , PubMed: <a href="#">12446774</a> , PubMed: <a href="#">14559997</a> , PubMed: <a href="#">14988723</a> , PubMed: <a href="#">15311285</a> , PubMed: <a href="#">15650047</a> , PubMed: <a href="#">15665856</a> , PubMed: <a href="#">32357935</a> ). May also negatively regulate cell cycle progression during unperturbed cell cycles (PubMed: <a href="#">11535615</a> , PubMed: <a href="#">12399544</a> , PubMed: <a href="#">12446774</a> , PubMed: <a href="#">14559997</a> , PubMed: <a href="#">14988723</a> , PubMed: <a href="#">15311285</a> , PubMed: <a href="#">15650047</a> , PubMed: <a href="#">15665856</a> ). This regulation is achieved by a number of mechanisms that together help to preserve the integrity of the genome (PubMed: <a href="#">11535615</a> , PubMed: <a href="#">12399544</a> , PubMed: <a href="#">12446774</a> , PubMed: <a href="#">14559997</a> , PubMed: <a href="#">14988723</a> , PubMed: <a href="#">15311285</a> , PubMed: <a href="#">15650047</a> , PubMed: <a href="#">15665856</a> ). Recognizes the |

substrate consensus sequence [R-X-X- S/T] (PubMed:[11535615](#), PubMed:[12399544](#), PubMed:[12446774](#), PubMed:[14559997](#), PubMed:[14988723](#), PubMed:[15311285](#), PubMed:[15650047](#), PubMed:[15665856](#)). Binds to and phosphorylates CDC25A, CDC25B and CDC25C (PubMed:[12676583](#), PubMed:[12676925](#), PubMed:[12759351](#), PubMed:[14559997](#), PubMed:[14681206](#), PubMed:[19734889](#), PubMed:[9278511](#)). Phosphorylation of CDC25A at 'Ser-178' and 'Thr-507' and phosphorylation of CDC25C at 'Ser-216' creates binding sites for 14-3-3 proteins which inhibit CDC25A and CDC25C (PubMed:[9278511](#)). Phosphorylation of CDC25A at 'Ser-76', 'Ser-124', 'Ser-178', 'Ser-279' and 'Ser-293' promotes proteolysis of CDC25A (PubMed:[12676583](#), PubMed:[12676925](#), PubMed:[12759351](#), PubMed:[14681206](#), PubMed:[19734889](#), PubMed:[9278511](#)). Phosphorylation of CDC25A at 'Ser-76' primes the protein for subsequent phosphorylation at 'Ser-79', 'Ser-82' and 'Ser-88' by NEK11, which is required for polyubiquitination and degradation of CDC25A (PubMed:[19734889](#), PubMed:[20090422](#), PubMed:[9278511](#)). Inhibition of CDC25 leads to increased inhibitory tyrosine phosphorylation of CDK-cyclin complexes and blocks cell cycle progression (PubMed:[9278511](#)). Also phosphorylates NEK6 (PubMed:[18728393](#)). Binds to and phosphorylates RAD51 at 'Thr-309', which promotes the release of RAD51 from BRCA2 and enhances the association of RAD51 with chromatin, thereby promoting DNA repair by homologous recombination (PubMed:[15665856](#)). Phosphorylates multiple sites within the C-terminus of TP53, which promotes activation of TP53 by acetylation and promotes cell cycle arrest and suppression of cellular proliferation (PubMed:[10673501](#), PubMed:[15659650](#), PubMed:[16511572](#)). Also promotes repair of DNA cross-links through phosphorylation of FANCE (PubMed:[17296736](#)). Binds to and phosphorylates TLK1 at 'Ser-743', which prevents the TLK1-dependent phosphorylation of the chromatin assembly factor ASF1A (PubMed:[12660173](#), PubMed:[12955071](#)). This may enhance chromatin assembly both in the presence or absence of DNA damage (PubMed:[12660173](#), PubMed:[12955071](#)). May also play a role in replication fork maintenance through regulation of PCNA (PubMed:[18451105](#)). May regulate the transcription of genes that regulate cell-cycle progression through the phosphorylation of histones (By similarity). Phosphorylates histone H3.1 (to form H3T11ph), which leads to epigenetic inhibition of a subset of genes (By similarity). May also phosphorylate RB1 to promote its interaction with the E2F family of transcription factors and subsequent cell cycle arrest (PubMed:[17380128](#)). Phosphorylates SPRTN, promoting SPRTN recruitment to chromatin (PubMed:[31316063](#)). Reduces replication stress and activates the G2/M checkpoint, by phosphorylating and inactivating PABIR1/FAM122A and promoting the serine/threonine-protein phosphatase 2A-mediated dephosphorylation and stabilization of WEE1 levels and activity (PubMed:[33108758](#)).

### Cellular Location

Nucleus. Chromosome. Cytoplasm Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Nuclear export is mediated at least in part by XPO1/CRM1 (PubMed:[12676962](#)). Also localizes to the centrosome specifically during interphase, where it may protect centrosomal CDC2 kinase from inappropriate activation by cytoplasmic CDC25B (PubMed:[15311285](#)). Proteolytic cleavage at the C-terminus by SPRTN promotes removal from chromatin (PubMed:[31316063](#))

### Tissue Location


Expressed ubiquitously with the most abundant expression in thymus, testis, small intestine and colon

## Background


The protein encoded by this gene belongs to the Ser/Thr protein kinase family. It is required for checkpoint mediated cell cycle arrest in response to DNA damage or the presence of unreplicated DNA. This protein

acts to integrate signals from ATM and ATR, two cell cycle proteins involved in DNA damage responses, that also associate with chromatin in meiotic prophase I. Phosphorylation of CDC25A protein phosphatase by this protein is required for cells to delay cell cycle progression in response to double-strand DNA breaks. Several alternatively spliced transcript variants have been found for this gene.



## Images



Western blot analysis of Chk1 in K562 lysates using Chk1 antibody.



Western blot analysis of Chk1 in K562, 3T3, HeLa lysates using Chk1 antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.