

# Cullin 2 Rabbit mAb

Catalog # AP75303

## Product Information

---

|                          |                        |
|--------------------------|------------------------|
| <b>Application</b>       | WB, FC, IP             |
| <b>Primary Accession</b> | <a href="#">Q13617</a> |
| <b>Reactivity</b>        | Rat, Human, Mouse      |
| <b>Host</b>              | Rabbit                 |
| <b>Clonality</b>         | Monoclonal Antibody    |
| <b>Isotype</b>           | IgG                    |
| <b>Conjugate</b>         | Unconjugated           |
| <b>Purification</b>      | Affinity Purified      |
| <b>Calculated MW</b>     | 87 KDa                 |

## Additional Information

---

|                    |   |
|--------------------|---|
| <b>Other Names</b> | CUL2  |
| <b>Dilution</b>    | WB~~1:1000-1:5000 FC~~1:50-1:100 IP~~1:10-1:100   |
| <b>Format</b>      | Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. |
| <b>Storage</b>     | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.        |

## Protein Information

---

### Background




---

Enables ubiquitin protein ligase binding activity. Predicted to be involved in SCF-dependent proteasomal ubiquitin-dependent protein catabolic process and protein ubiquitination. Predicted to act upstream of or within protein catabolic process. Located in nucleoplasm. Part of Cul2-RING ubiquitin ligase complex.

### Images

---

Western blot analysis of Cullin 2/CUL2 in K562, C6, 3T3, Hela lysates using Cullin 2 antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.